arXiv:1004.5246v1 [physics.atom-ph] 29 Apr 2010

Irreducible tensor form of three-particle operator for open-shell atoms

R. Juršėnas and G. Merkelis

Institute of Theoretical Physics and Astronomy of Vilnius University, A. Goštauto 12, LT-01108,

Abstract

The three-particle operator in a second quantized form is studied. The operator is transformed into irreducible tensor form. Possible coupling schemes, distinguished by the classes of symmetric group S_6, are presented. Recoupling coefficients, which allow one to transform given scheme into another, are produced by using the angular momentum theory, combined with quasispin formalism. The classification of three-particle operator, which acts on $n = 1, 2, \ldots, 6$ open shells of equivalent electrons of atom, is considered. The procedure to construct three-particle matrix elements are examined.

Keywords: Three-particle operator, Tensor operator, Recoupling coefficients, Quasispin

1 Introduction

The mathematical technique of effective operators plays an important role in the studies of either atomic open-shell many-body perturbation theories [1–4] or atomic inner-shell physics, when calculating transition probabilities or widths of levels [5, 6]. This approach provides an opportunity to include electron-electron correlation effects in a flexible form. The irreducible tensor forms of one-, two-particle effective operators have been widely studied in many works [7–11]. Starting from the second-order MBPT, in a particle-hole representation one has to deal with expansion terms, which include three-particle operators. In this paper we namely concentrate on classification of effective three-particle operator, written in irreducible tensor form. Special attention is paid for many open-shell aspects of the problem. The group-theoretical classification of irreducible three-particle operator for equivalent electrons (particularly, f electrons) has been considered by Judd [7]. In our considerations we do not classify the states of equivalent electrons. Our aim is the operator by itself, providing a general classification of all possible coupling schemes and determining connection among them.

The three-particle operator may inscribe on six open-shell electrons at once. It is assumed that irreducible tensor operator matrix element is constructed on the basis of multi-shell wave functions, coupled in a consequent order. For this reason, tensor operator is also expressed in a consequent order with respect to creation and annihilation operators, which act on the first, second, etc. open-shell. All other mixed arrangements are obtained by the corresponding permutation operations of S_6 group. Recoupling coefficients, which appear due to such transformations, are presented too.

Besides the tensor structure, effective three-particle operator includes weight coefficient. In MBPT this coefficient denotes miscellaneous products of one- or two-particle matrix elements (with energy denominator included). The systematic study of two-particle matrix elements can be found in [12–14]. The application of this methodology for the second-order effective Hamiltonian one can find in [15].

2 Preliminaries

Let a generalized three-particle operator be

$$\widehat{L}_3 = \sum_{ijk}\sum_{lpq} a_i a_j a_k a_l^{\dagger} a_p^{\dagger} a_q^{\dagger}\ \ell_{ijkqpl}. \tag{2.1}$$

where ℓ_{ijkqpl} denotes weight coefficient, which depends on concrete operator. The subscripts $x = i, j, \ldots$ indicate the sets of one-electron quantum numbers $\{n_x \lambda_x \mu_x\}$. Let Λ be either reducible (LS-coupling) or irreducible (jj-coupling) tensor space. Then each creation operator a_x is labeled by irreducible representations $\lambda_x = l_x \frac{1}{2}$ (in LS-coupling) or $\lambda_x = j_x$ (in jj-coupling), denoting $a^{\lambda_x}_{\mu_x}$. The reduction of Eq. (2.1) is performed transposing annihilation operator by $\tilde{a}^{\lambda_x}_{\mu_x} = (-1)^{\lambda_x - \mu_x} a^{\lambda_x \dagger}_{-\mu_x}$, and exploiting known reduction rules for the Kronecker product $\lambda_x \times \lambda_{\bar{x}} \rightarrow \lambda_{x\bar{x}}$ of SU(2)-irreducible representations. For instance,

$$\widehat{L}_3 = \sum_{n_x \lambda_x \mu_x} \sum_{\lambda_{ij} \lambda_{ijk}} \sum_{\lambda_{lp} \lambda_{lpq}} \sum_{\lambda \mu} (-1)^{\lambda_l + \lambda_p + \lambda_q + \mu_l + \mu_p + \mu_q} \ell_{ijkqpl} I \ {}_{\Lambda}\widehat{T}^{\lambda}_{\mu} \left(\lambda_i \lambda_j \lambda_k \lambda_l \lambda_p \lambda_q\right), \tag{2.2}$$

$$I \equiv I \left[\lambda_i \lambda_j \left(\lambda_{ij}\right) \lambda_k \left(\lambda_{ijk}\right), \lambda_l \lambda_p \left(\lambda_{lp}\right) \lambda_q \left(\lambda_{lpq}\right) \lambda \mu \left\{\mu_x\right\}\right]$$

$$= \sum_{\mu_{ij} \mu_{ijk}} \sum_{\mu_{lp} \mu_{lpq}} \left\langle \lambda_i \mu_i \lambda_j \mu_j | \lambda_{ij} \mu_{ij} \right\rangle \left\langle \lambda_{ij} \mu_{ij} \lambda_k \mu_k | \lambda_{ijk} \mu_{ijk} \right\rangle \left\langle \lambda_l \mu_l \lambda_p \mu_p | \lambda_{lp} \mu_{lp} \right\rangle$$

$$\times \left\langle \lambda_{lp} \mu_{lp} \lambda_q \mu_q | \lambda_{lpq} \mu_{lpq} \right\rangle \left\langle \lambda_{ijk} \mu_{ijk} \lambda_{lpq} \mu_{lpq} | \lambda \mu \right\rangle, \tag{2.3}$$

$${}_{\Lambda}\widehat{T}^{\lambda}_{\mu} \left(\lambda_i \lambda_j \lambda_k \lambda_l \lambda_p \lambda_q\right) = \left[\left[\left[a^{\lambda_i} \times a^{\lambda_j}\right]^{\lambda_{ij}} \times a^{\lambda_k}\right]^{\lambda_{ijk}} \times \left[\left[\tilde{a}^{\lambda_l} \times \tilde{a}^{\lambda_p}\right]^{\lambda_{lp}} \times \tilde{a}^{\lambda_q}\right]^{\lambda_{lpq}}\right]^{\lambda}_{\mu}. \tag{2.4}$$

In Eq. (2.3) the quantities $\left\langle \lambda_x \mu_x \lambda_y \mu_y | \lambda_{xy} \mu_{xy} \right\rangle$ denote Clebsch-Gordan coefficients of SU (2) [16]. Set Q a quasispin space. Then the presentation of ${}_{\Lambda}\widehat{T}^{\lambda}_{\mu}$ in q-space, where $q \equiv Q \times \Lambda$, is

$$\widehat{T}^{\varsigma}_{\iota} \left(\lambda_i \lambda_j \lambda_k \lambda_l \lambda_p \lambda_q\right) = \langle ijklpq \rangle = [[[a^{\varsigma_i} \times a^{\varsigma_j}]^{\varsigma_{ij}} \times a^{\varsigma_k}]^{\varsigma_{ijk}} \times [[a^{\varsigma_l} \times a^{\varsigma_p}]^{\varsigma_{lp}} \times a^{\varsigma_q}]^{\varsigma_{lpq}}]^{\varsigma}_{\iota}. \tag{2.5}$$

The notation $\langle ijklpq \rangle$, as a shorter description of $\widehat{T}^{\varsigma}_{\iota}$, will be used later. The ranks $\varsigma_x = \kappa_x \lambda_x$. In the next section it will be showed, that there are 42 ways at all to form $\widehat{T}^{\varsigma}_{\iota}$.

3 Coupling schemes

Suppose there is constructed tensor

$$\widehat{O}_n = T^{\alpha_1 \alpha_2 \ldots \alpha_n}_{\beta_1 \beta_2 \ldots \beta_n} = a^{\alpha_1}_{\beta_1} a^{\alpha_2}_{\beta_2} \ldots a^{\alpha_n}_{\beta_n} \tag{3.1}$$

with $n \geq 2$, and α_i, $i \in \mathcal{I} = \{1, 2, \ldots, 6\}$ labeling irreducible representations. In general, for any $n \geq 2$ there are different ways to reduce Kronecker products of α_i. We suggest a procedure, suitable for any n, to obtain possible coupling schemes. Suppose there is a n-length string, which is partitioned into the sum of $\lambda_{\varkappa}$-length strings, where $n = \lambda_1 + \lambda_2 + \ldots + \lambda_n$. Consequently, the partitions can be characterized by the irreducible representations $\lambda \equiv [\lambda_1, \lambda_2, \ldots \lambda_{\varkappa}, \ldots, \lambda_n]$ of S_n. We exclude two partitions of length 1 and 2. The 1-partition, including only 1 operator $a^{\alpha_i}_{\beta_i}$, appears h_1 times in the coupling scheme, the 2-partition appears h_2 times in the same coupling scheme and so on. Furthermore, in our considerations the ordering of partitions is important, while the ordering of partitions $\lambda_{\varkappa}$ in the sum $\lambda_1 + \lambda_2 + \ldots + \lambda_n$ does not play any role[1]. In order to classify irreducible tensor operators by coupling schemes, we use n-tuples which, by default, represent the notation of an ordered list of elements in the scheme. Such elements will be one- and two-operator partitions, denoted by the numbers 1 and 2. For example, the notation of the scheme $[\bullet \times \bullet]$ is $[[2]]$; the schemes $[[\bullet \times \bullet] \times \bullet]$ and $[\bullet \times [\bullet \times \bullet]]$ are labeled by $[[21]]$ and $[[12]]$, respectively. Here it is assumed that in the bullets $\bullet$ there are placed the operators $a^{\alpha_i}_{\beta_i}$ and the subscript i marks the position of given operator (or bullet) starting from the left. One may notice, the 3-tuples $[[21]]$ and $[[12]]$ are characterized by the same irreducible representation $[21]$, thus they belong to the same class $(\alpha) = (1^1 2^1)$ of S_3.

---

[1] By definition, $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_n$. However, in our case $\lambda_i + \lambda_j$ has different meaning than $\lambda_j + \lambda_i$.

(α)	Tuples	Symbol
(2^1)	$[[2]]$	$\widehat{\mathrm{T}}^{[1^2]}$
$(1^1 2^1)$	$[[12]]$	$\widehat{\mathrm{T}}^{[21]}_{1}$
	$[[21]]$	$\widehat{\mathrm{T}}^{[21]}_{2}$
(2^2)	$[[22]]$	$\widehat{\mathrm{T}}^{[2^2]}$
$(1^1 3^1)$	$[[211]] \ltimes \widehat{\mathrm{T}}^{[21]}_{2}$	$\widehat{\mathrm{T}}^{[21^2]}_{1}$
	$[[121]] \ltimes \widehat{\mathrm{T}}^{[21]}_{2,1}$	$\widehat{\mathrm{T}}^{[21^2]}_{2,3}$
	$[[112]] \ltimes \widehat{\mathrm{T}}^{[21]}_{1}$	$\widehat{\mathrm{T}}^{[21^2]}_{4}$
$(2^1 3^1)$	$[[221]] \ltimes \widehat{\mathrm{T}}^{[21]}_{2,1}$	$\widehat{\mathrm{T}}^{[2^2 1]}_{1,2}$
	$[[122]] \ltimes \widehat{\mathrm{T}}^{[21]}_{1,2}$	$\widehat{\mathrm{T}}^{[2^2 1]}_{3,4}$
	$[[212]] \ltimes \widehat{\mathrm{T}}^{[21]}_{2,1}$	$\widehat{\mathrm{T}}^{[2^2 1]}_{5,6}$
$(1^1 4^1)$	$[[2111]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1}$	$\widehat{\mathrm{T}}^{[21^3]}_{1}$
	$[[1211]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1,3,2}$	$\widehat{\mathrm{T}}^{[21^3]}_{2,3,4}$
	$[[1121]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{4,2,3}$	$\widehat{\mathrm{T}}^{[21^3]}_{5,6,7}$
	$[[1112]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{4}$	$\widehat{\mathrm{T}}^{[21^3]}_{8}$

Tab. 1: The coupling schemes for $\widehat{O}_{1-5}$

(α)	Tuples	Symbol
(3^2)	$[[222]] \ltimes \widehat{\mathrm{T}}^{[21]}_{2,1}$	$\widehat{\mathrm{T}}^{[2^3]}_{1,2}$
$(2^1 4^1)$	$[[2211]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1,2,3}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{1,2,3}$
	$[[1221]] \ltimes \widehat{\mathrm{T}}^{[2^2]}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{4}$
	$[[1221]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1-4}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{5-8}$
	$[[1122]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{2,3,4}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{9,10,11}$
	$[[2121]] \ltimes \widehat{\mathrm{T}}^{[2^2]}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{12}$
	$[[2121]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1-4}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{13-16}$
	$[[2112]] \ltimes \widehat{\mathrm{T}}^{[2^2]}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{17}$
	$[[2112]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1,4}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{18,19}$
	$[[1212]] \ltimes \widehat{\mathrm{T}}^{[2^2]}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{20}$
	$[[1212]] \ltimes \widehat{\mathrm{T}}^{[21^2]}_{1-4}$	$\widehat{\mathrm{T}}^{[2^2 1^2]}_{21-24}$
$(1^1 5^1)$	$[[21111]] \ltimes \widehat{\mathrm{T}}^{[21^3]}_{1}$	$\widehat{\mathrm{T}}^{[21^4]}_{1}$
	$[[12111]] \ltimes \widehat{\mathrm{T}}^{[21^3]}_{1-4}$	$\widehat{\mathrm{T}}^{[21^4]}_{2-5}$
	$[[11211]] \ltimes \widehat{\mathrm{T}}^{[21^3]}_{2-7}$	$\widehat{\mathrm{T}}^{[21^4]}_{6-11}$
	$[[11121]] \ltimes \widehat{\mathrm{T}}^{[21^3]}_{5-8}$	$\widehat{\mathrm{T}}^{[21^4]}_{12-15}$
	$[[11112]] \ltimes \widehat{\mathrm{T}}^{[21^3]}_{8}$	$\widehat{\mathrm{T}}^{[21^4]}_{16}$

Tab. 2: The coupling schemes for $\widehat{O}_6$

Suppose $n = 2$. Then $\lambda_1 = [2]$ and $\lambda_2 = [1^2]$. We distinguish the antisymmetric S_2-irreducible representation. The tuple $[[2]]$ is denoted by $\widehat{\mathrm{T}}^{[1^2]}$ (see Tab. 1). If $n = 3$, S_3-irreducible representation, conforming to 1- and 2-partitions, is $\lambda = [21]$ (the full set of partitions equals to $[3]$, $[21]$, $[1^3]$). It corresponds to Kronecker products $(\alpha_1 \times \alpha_2) \times \alpha_3$ or $\alpha_1 \times (\alpha_2 \times \alpha_3)$ (previously studied example of 3-tuples $[[21]]$, $[[12]]$). Further, when $n = 4$, the suitable representations are $[2^2]$ and $[21^2]$. Let us study the schemes of class $(1^1 3^1)$. Suppose there is a tuple $\widehat{\mathrm{T}}^{[21^2]}_{1} = [[211]] \ltimes \widehat{\mathrm{T}}^{[21]}_{2}$, where $\widehat{\mathrm{T}}^{[21]}_{2} = [[21]]$ (Tab. 1). The tuple $\widehat{\mathrm{T}}^{[21]}_{2}$ in an explicit form reads $[[\bullet \times \bullet] \times \bullet]$. The (semijoin) notation $\ltimes$ denotes that the scheme $\widehat{\mathrm{T}}^{[21^2]}_{1}$ consists of only those partitions, which are in the scheme $\widehat{\mathrm{T}}^{[21]}_{2}$. The tuple $[[211]]$ reads $[\bullet \times \bullet] \times \bullet \times \bullet$. If we rename $[\bullet \times \bullet] = \bullet'$, then it becomes of the form $\bullet' \times \bullet \times \bullet$. We obtained the case of $n = 3$. Then, according to $\widehat{\mathrm{T}}^{[21]}_{2}$, the form $\bullet' \times \bullet \times \bullet$ is partitioned as $[[\bullet' \times \bullet] \times \bullet]$. Thus, the final scheme $\widehat{\mathrm{T}}^{[21^2]}_{1}$ reads $[[[\bullet \times \bullet] \times \bullet] \times \bullet]$.

Analogously possible schemes are obtained for higher n. When $n = 5$, the suitable representations of S_5 are $[2^2 1]$ and $[21^3]$. Finally, when $n = 6$, the representations are $[2^3]$, $[2^2 1^2]$ and $[21^4]$ (see Tab. 2). Particularly, the tuple $\widehat{\mathrm{T}}^{[2^2 1^2]}_{12}$ corresponds to coupling scheme of ranks ς_x in $\widehat{T}^{\varsigma}$ in Eq. (2.5). In Tabs. 1-2 notations $\widehat{\mathrm{T}}^{\lambda}_{a,b,c,\ldots} = [[x]] \ltimes \widehat{\mathrm{T}}^{\lambda'}_{a',b',c',\ldots}$ signify that $\widehat{\mathrm{T}}^{\lambda}_{a} = [[x]] \ltimes \widehat{\mathrm{T}}^{\lambda'}_{a'}$, $\widehat{\mathrm{T}}^{\lambda}_{b} = [[x]] \ltimes \widehat{\mathrm{T}}^{\lambda'}_{b'}$ and so on. The procedure considered is easily extended for any n, if there are defined all m-tuples with $m < n$. To perform this task, the following steps should be accomplished: (i) each n-length string is distinguished by representations of S_n, excluding only those, which include 1- and 2-partitions; (ii) obtained partitions $\lambda_{\varkappa}$ are written in all possible ways in the sum $\lambda_1 + \lambda_2 + \ldots + \lambda_n$ (e.g., $2 + 1 + 1$, $1 + 2 + 1$, $1 + 1 + 2$); (iii) renaming 2 by $1'$, one obtains $(n - h_2)$-length string (h_2 is a multiplicity of 2), which has already been classified by possible partitions; those partitions are included in studied string (by using $\ltimes$), and in the final step $1'$ is reversed to 2 back again. Note, for obtained $(n-h_2)$-length string there are suitable only those partitions, which do not couple 1-partitions; otherwise one would get a tuple, characterized by another representation of S_n (e.g., $[[1'\times 1]\times 1]$, $[[1\times 1']\times 1]$ or $[1\times[1'\times 1]]$, $[1\times[1\times 1']]$, but not $[1'\times[1\times 1]]$, $[[1\times 1]\times 1']$, which correspond to $\lambda=[2^2]$, rather than $\lambda=[21^2]$). Finally, substituting in the bullets operators a^{α_i}, and each 2-partition renaming by tensor operator $W^{\alpha_{ij}}(\lambda_i\lambda_j)$, where $W^{\alpha_{ij}}(\lambda_i\lambda_j)=[a^{\alpha_i}\times a^{\alpha_j}]^{\alpha_{ij}}$, we would gain irreducible tensor operators for given reducible product $\widehat{O}_n$.

In general, there are $42\cdot 42=1764$ recoupling coefficients which can be generated among each of the scheme (written in a consequent order) when $n=6$ (Tab. 2), including coefficients for the same scheme and for the conjugate schemes. This number can be reduced to only 42. We will call them the basis coefficients. All other recoupling coefficients are generated from basis coefficients. Let it be the coefficients generated when transforming $\widehat{T}^{\varsigma}_{\iota}\equiv\widehat{\mathrm{T}}^{[2^21^2]}_{12}$ into all other 41 scheme. Here

$$\widehat{\mathrm{T}}^{[2^21^2]}_{12}\equiv[[[a^{\alpha_1}\times a^{\alpha_2}]^{\alpha_{12}}\times a^{\alpha_3}]^{\alpha_{123}}\times[[a^{\alpha_4}\times a^{\alpha_5}]^{\alpha_{45}}\times a^{\alpha_6}]^{\alpha_{456}}]^{\alpha}_{\beta}\,. \tag{3.2}$$

Then the tensor $\widehat{O}_6$ (see Eq. (3.1)) is reduced as follows

$$\widehat{O}_6=T^{\alpha_1\alpha_2\ldots\alpha_6}_{\beta_1\beta_2\ldots\beta_6}=\underset{\alpha_{12}\alpha_{123}\alpha_{45}\alpha_{456}}{\sum\sum}\widehat{\mathrm{T}}^{[2^21^2]}_{12}\,I. \tag{3.3}$$

The multiplier I is defined in Eq. (2.3) making the replacements $\lambda_x\mu_x\to\varsigma_x\iota_x$, where $\varsigma_i=\alpha_1$, $\varsigma_j=\alpha_2$, $\varsigma_k=\alpha_3$, $\varsigma_l=\alpha_4$, $\varsigma_p=\alpha_5$, $\varsigma_q=\alpha_6$, $\varsigma=\alpha_{123456}\equiv\alpha$, $\iota=\beta$. It has to be taken into account that the numbering of the ranks and of the operators differs. For example, the rank α_{12} is the total rank of the ranks α_1 and α_2 of the operators which are placed in the 1st and 2nd positions of the given scheme. On the other hand, the ranks $\varsigma_i=\alpha_1$ and $\varsigma_j=\alpha_2$ can acquire the same value ς_1. To avoid further misunderstanding, the following notations will be used. The rank $\alpha_i=\kappa_i\lambda_i$ will denote the rank of the ith operator a^{α_i} in q-space; the rank $\varsigma_x=\kappa_x\lambda_x$ will denote some value of any of the operators in the sequence. Generally, if the notations α_i and α_j mark the ith and the jth positions of the corresponding operators, respectively, the rank ς_x can be the same for both α_i and α_j.

The basis coefficients will be denoted by ϵ_ξ, where

$$\epsilon_\xi=\sum_{\beta_i\in\mathcal{I}}I\cdot I_\xi. \tag{3.4}$$

The integers $\xi\in\{1,2,\ldots,42\}$; $I_\xi=I_\varkappa=I^{[2^3]}_\varkappa$ for $\varkappa=1,2$ and $\lambda=[2^3]$; $I_\xi=I_{\varkappa+2}=I^{[2^21^2]}_\varkappa$ for $\varkappa\in[1,24]$ and $\lambda=[2^21^2]$; $I_\xi=I_{\varkappa+24}=I^{[21^4]}_\varkappa$ for $\varkappa\in[1,16]$ and $\lambda=[21^4]$. Then follows $I_{14}=I$. The coefficient $I^\lambda_\varkappa$, in general, depends on the partition $\widehat{\mathrm{T}}^\lambda_\varkappa$ in the following way

$$\widehat{\mathrm{T}}^\lambda_\varkappa=\sum_{\beta_i\in\mathcal{I}}\widehat{O}_6\,I^\lambda_\varkappa. \tag{3.5}$$

According to Eqs. (3.3), (3.5), the schemes $\widehat{\mathrm{T}}^\lambda_\varkappa$ and $\widehat{\mathrm{T}}^{[2^21^2]}_{12}$ are transforming among each other by the formulas

$$\widehat{\mathrm{T}}^\lambda_\varkappa=\sum_{\alpha_\eta\in\Upsilon}\widehat{\mathrm{T}}^{[2^21^2]}_{12}\,\epsilon_\xi,\quad \widehat{\mathrm{T}}^{[2^21^2]}_{12}=\sum_{\alpha_\zeta\in\Gamma_\xi}\widehat{\mathrm{T}}^\lambda_\varkappa\,\epsilon_\xi, \tag{3.6}$$

where the summations are performed over the internal ranks α_η, α_ζ. The indices $\eta\in\Upsilon=\{12,123,45,456\}$; the indices $\zeta\in\Gamma_\xi=\{\zeta_1,\zeta_2,\zeta_3,\zeta_4\}$ depend on the concrete scheme $\widehat{\mathrm{T}}^\lambda_\varkappa$. It directly follows from Eq. (3.6) that

$$\widehat{\mathrm{T}}^\lambda_\varkappa=\sum_{\alpha_\eta\in\Upsilon}\sum_{\alpha_{\zeta'}\in\Gamma_{\xi'}}\widehat{\mathrm{T}}^{\lambda'}_{\varkappa'}\,\epsilon_\xi\epsilon_{\xi'},\quad \sum_{\alpha_\eta\in\Upsilon}\epsilon^2_\xi=1. \tag{3.7}$$

In Eq. (3.7) the basis coefficients ϵ_{ξ} are defined in Eq. (3.4), while the basis coefficients $\epsilon_{\xi'}$ correspond to the recoupling from $\widehat{\mathrm{T}}_{12}^{[2^2 1^2]}$ to $\widehat{\mathrm{T}}_{\varkappa'}^{\lambda'}$. The coefficients ϵ_{ξ} are presented in an explicit form in Appendix A.

4 Permutations

In previous section the possible schemes to reduce operator $\widehat{O}_6$ have been studied. In present section we develop the procedure of transformation for arbitrary operator $\widehat{T}^{\varsigma}(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q)$ with i, j, k, l, p, q labeling open shells, into $\widehat{T}^{\varsigma}(\lambda_{i'}\lambda_{j'}\lambda_{k'}\lambda_{l'}\lambda_{p'}\lambda_{q'})$, where $i' \leq j' \leq k' \leq l' \leq p' \leq q'$. It is assumed, both operators are coupled by the scheme $\widehat{\mathrm{T}}_{12}^{[2^2 1^2]}$ (see Eqs. (2.5), (3.2)). Recoupling coefficients, which appear due to the permutations of ranks of $T_{\beta_1\ldots\beta_6}^{\alpha_1\ldots\alpha_6}$, written in irreducible tensor form $\widehat{T}_{\beta}^{\alpha}$, will be called the permutation coefficients, denoted ε. It is clear, there are 6! possible permutations for given distributions i, j, k, l, p, q, if $i \neq j \neq k \neq l \neq p \neq q$. Consequently, there are 6! permutation coefficients ε, in general. This number will be reduced up to 15 permutations which are described by the 2-cycles $(ij) \in \mathrm{S}_6$ $(i, j \in \mathcal{I})$. This appears from the general theory which states that all n-cycles $(n \geq 2)$ can be expanded into the composition of $n-1$ number of 2-cycles. Hence, the rest of permutation coefficients are obtained from the permutation coefficients ε_{ij}. Moreover, the coefficients ε_{ij} will be constructed of the basis coefficients ϵ_{ξ} in such a way, that the desirable ordering of ranks of the tensor operator $\widehat{T}_{\beta}^{\alpha}$ would be obtained making minimal number of recouplings between the schemes $\widehat{\mathrm{T}}_{\varkappa}^{\lambda}$. The coefficients ε_{ij} are determined from the equation[2]

$$(ij)\, \widehat{\mathrm{T}}_{12}^{[2^2 1^2]} = \sum_{\alpha_{\varrho \in \mathcal{F}}} \varepsilon_{ij} \widehat{\mathrm{T}}_{12}^{[2^2 1^2]}. \tag{4.1}$$

The summation is performed over α_{ϱ}, where the indices $\varrho \in \mathcal{F} \subset \mathcal{H} = \{12, 123, 45, 46, 456\}$ depend on the concrete permutation (ij). These coefficients ε_{ij} are invariant under the operation (ij)

$$(ij)\, \varepsilon_{ij} = \varepsilon_{ij}. \tag{4.2}$$

However, in general, $(ij)\, \varepsilon_{ik} \neq \varepsilon_{jk}$. Some of the coefficients $(ij)\, \varepsilon_{ik}$ and ε_{jk} differ only by the phase factor

$$\begin{cases} (12)\, \varepsilon_{1k} = f_{1k} f_{2k} \varepsilon_{2k}, \quad k \in \{3, 4, 5, 6\}, \\ (45)\, \varepsilon_{34} = f_{34} f_{35} \varepsilon_{35}, \\ (45)\, \varepsilon_{46} = f_{46} f_{56} \varepsilon_{56}. \end{cases} \tag{4.3}$$

Here

$$f_{ij} = \prod_{k=1}^{j-i} \prod_{l=1}^{j-i-1} \varpi_{i\, i+k}\, \varpi_{i+l\, j} \qquad \forall j > i; \quad \varpi_{ii} = 1, \quad \varpi_{ij} = (-1)^{\alpha_i + \alpha_j - \alpha_{ij} + 1}. \tag{4.4}$$

Since $\varpi_{ij} = \varpi_{ji} = \varpi_{ij}^{-1}$, the condition $j > i$ can always be satisfied. We conclude that any operation of the n-length $(2 \leq n \leq 6)$ permutation $(i_1 i_2 \ldots i_n)$ on the scheme $\widehat{\mathrm{T}}_{12}^{[2^2 1^2]}$ can be decomposed into the $n-1$ power sums of the operations of 2-cycles (ij) on given scheme by applying Eq. (4.1) in the following way

$$\begin{aligned} (ij) \ldots (kl)\, (pq)\, \widehat{\mathrm{T}}_{12}^{[2^2 1^2]} &= (ij) \ldots (kl) \sum_{\alpha_{\varrho \in \mathcal{F}}} \varepsilon_{pq} \widehat{\mathrm{T}}_{12}^{[2^2 1^2]} = (ij) \ldots \sum_{\alpha_{\varrho \in (kl)\mathcal{F}}} \sum_{\alpha_{\varrho' \in \mathcal{F}'}} \varepsilon_{kl} \left\{ (kl)\, \varepsilon_{pq} \right\} \widehat{\mathrm{T}}_{12}^{[2^2 1^2]} \\ &= (ij) \sum_{\alpha_{\varrho \in \ldots (kl)\mathcal{F}}} \sum_{\alpha_{\varrho' \in \ldots \mathcal{F}'}} \cdots \sum_{\alpha_{\varrho'' \in \mathcal{F}''}} \ldots \left\{ \ldots \varepsilon_{kl} \right\} \left\{ \ldots (kl)\, \varepsilon_{pq} \right\} \widehat{\mathrm{T}}_{12}^{[2^2 1^2]} \\ &= \sum_{\alpha_{\varrho \in (ij) \ldots (kl)\mathcal{F}}} \sum_{\alpha_{\varrho' \in (ij) \ldots \mathcal{F}'}} \cdots \sum_{\alpha_{\varrho'' \in \ldots \mathcal{F}''}} \sum_{\alpha_{\varrho''' \in \mathcal{F}'''}} \varepsilon_{ij} \ldots \left\{ (ij) \ldots \varepsilon_{kl} \right\} \left\{ (ij) \ldots (kl)\, \varepsilon_{pq} \right\} \widehat{\mathrm{T}}_{12}^{[2^2 1^2]}. \end{aligned} \tag{4.5}$$

---

[2] Hereafter we formally distinguish permutation operators $\widehat{\pi}$ on n-particle Hilbert space and elements π of S_n. However, a cycle notation $(ij \ldots k)$ suits for both cases.

In Tab. 3 the explicit forms of ε_{ij} are presented for all 15 cases. In the last column the set $\mathcal{F}$ of α_{ϱ} indices is defined in Eq. (4.1). The number $\xi > 1$ signifies, that there are more than one equivalent opportunities to recouple given operator $\widehat{T}^{\varsigma}(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q)$ into $\widehat{T}^{\varsigma}(\lambda_{i'}\lambda_{j'}\lambda_{k'}\lambda_{l'}\lambda_{p'}\lambda_{q'})$. By default, the coefficients ε_{ij} appear when permuting the operators $a_{\iota_x}^{\varsigma_x}$ $(x = i, j, k, l, p, q \in \mathcal{I})$ of $\widehat{\mathrm{T}}_{12}^{[2^2 1^2]}$ with different ranks ς_x. It is noticeable, some of the indices x can acquire the same values, i.e., some ranks can be equal. In this case one assumes that the same rank operators $a_{\iota_x}^{\varsigma_x}$ are not permuted because in a contrary case the Kronecker delta function appears. However, we will demonstrate, that there is a simple way to avoid arising deltas. Suppose there is a tensor operator $\langle jiilpq\rangle$ (see Eq. (2.5)) which has to be transformed to $\langle iijlpq\rangle$. The set of possible operators $\widehat{\pi}$ satisfying such a transformation equals to $\{(13)\,,(132)\}$. If $\widehat{\pi} = (13) = \left(\begin{smallmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{smallmatrix}\right)$, then $(13)\,\langle iijlpq\rangle = \langle jiilpq\rangle = \sum_{\varsigma_{ii}=\mathrm{odd}} \varepsilon_{13}\,\langle iijlpq\rangle$, where $\varepsilon_{13} = f_{13}\,\{(12)\,\epsilon_6\}\,\{(123)\,\epsilon_6\}$, $f_{13} = \varpi_{12}\varpi_{13}\varpi_{23} = 1$, since $\varpi_{12} = 1$, $\varpi_{13} = \varpi_{23} = (-1)^{\varsigma_i+\varsigma_j+\varsigma_{ij}+1}$, and ϵ_6 is presented in Eq. (A.11).

(ij)	ε_{ij}	ξ	Γ_ξ	$\mathcal{F}$
(12)	f_{12}	1	$\emptyset$	$\emptyset$
(23)	$\sum_{\alpha_{23}} f_{23}\epsilon_6 \left\{(23)\,\epsilon_6\right\}$	1	$\emptyset$	$\{12\}$
(34)	$\sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{34}\epsilon_\xi \left\{(34)\,\epsilon_\xi\right\}$	1 2 3 4 5	$\{34,1234,56\}$ $\{34,56,3456\}$ $\{34,1234,12345\}$ $\{34,345,12345\}$ $\{34,345,3456\}$	$\{123,45\}$
(45)	f_{45}	1	$\emptyset$	$\emptyset$
(56)	$\sum_{\alpha_{56}} f_{56}\epsilon_{19} \left\{(56)\,\epsilon_{19}\right\}$	1	$\emptyset$	$\{45\}$
(13)	$\sum_{\alpha_{13}} f_{13} \left\{(12)\,\epsilon_6\right\} \left\{(123)\,\epsilon_6\right\}$	1	$\emptyset$	$\{12\}$
(24)	$\sum_{\alpha_{13}\alpha_{23}\alpha_{34}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{24}\epsilon_6 \left\{(24)\,\epsilon_6\right\}$ $\times \left\{(23)\,[\epsilon_6\epsilon_\xi]\right\} \left\{(234)\,[\epsilon_6\epsilon_\xi]\right\}$	1 2 3 4 5	$\{24,1234,56\}$ $\{24,56,2456\}$ $\{24,1234,12345\}$ $\{24,245,12345\}$ $\{24,245,2456\}$	$\{12,123,45,456\}$
(35)	$\sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{35} \left\{(45)\,\epsilon_\xi\right\} \left\{(354)\,\epsilon_\xi\right\}$	1 2 3 4 5	$\{35,46,1235\}$ $\{35,46,3456\}$ $\{35,1235,12345\}$ $\{35,345,12345\}$ $\{35,345,3456\}$	$\{123,45,456\}$
(46)	$\sum_{\alpha_{46}} f_{46} \left\{(45)\,\epsilon_{19}\right\} \left\{(456)\,\epsilon_{19}\right\}$	1	$\emptyset$	$\{45\}$
(14)	$\sum_{\alpha_{13}\alpha_{23}\alpha_{34}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{14} \left\{(12)\,\epsilon_6\right\} \left\{(124)\,\epsilon_6\right\}$ $\times \left\{(123)\,[\epsilon_6\epsilon_\xi]\right\} \left\{(1234)\,[\epsilon_6\epsilon_\xi]\right\}$	1 2 3 4 5	$\{14,1234,56\}$ $\{14,56,1456\}$ $\{14,1234,12345\}$ $\{14,145,12345\}$ $\{14,145,1456\}$	$\{12,123,45,456\}$
(25)	$\sum_{\alpha_{13}\alpha_{23}\alpha_{35}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{25}\epsilon_6 \left\{(254)\,\epsilon_6\right\}$ $\times \left\{(2354)\,[\epsilon_6\epsilon_\xi]\right\} \left\{(23)\,(45)\,[\epsilon_6\epsilon_\xi]\right\}$	1 2 3 4 5	$\{25,46,1235\}$ $\{25,46,2456\}$ $\{25,1235,12345\}$ $\{25,245,12345\}$ $\{25,245,2456\}$	$\{12,123,45,456\}$
(36)	$\sum_{\alpha_{34}\alpha_{35}\alpha_{56}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{36}\epsilon_{19} \left\{(36)\,\epsilon_{19}\right\} \left\{(56)\,\epsilon_{19}\right\}$ $\times \left\{(365)\,\epsilon_{19}\right\} \left\{(465)\,\epsilon_1\right\} \left\{(3654)\,\epsilon_1\right\}$	1 2	$\{36,1236\}$ $\{36,3456\}$	$\{123,46,456\}$
(15)	$\sum_{\alpha_{13}\alpha_{23}\alpha_{35}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{15} \left\{(12)\,\epsilon_6\right\} \left\{(1254)\,\epsilon_6\right\}$ $\times \left\{(12354)\,[\epsilon_6\epsilon_\xi]\right\} \left\{(123)\,(45)\,[\epsilon_6\epsilon_\xi]\right\}$	1 2 3 4 5	$\{15,46,1235\}$ $\{15,46,1456\}$ $\{15,1235,12345\}$ $\{15,145,12345\}$ $\{15,145,1456\}$	$\{12,123,45,456\}$
(26)	$\sum_{\alpha_{13}\alpha_{23}\alpha_{36}} \sum_{\alpha_{24}\alpha_{25}\alpha_{56}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{26}\epsilon_6$ $\times \left\{(23)\,[\epsilon_6\epsilon_{19}]\right\} \left\{(26)\,\epsilon_6\right\} \left\{(236)\,[\epsilon_6\epsilon_{19}]\right\}$ $\times \left\{(23)\,(56)\,\epsilon_{19}\right\} \left\{(2365)\,\epsilon_{19}\right\}$ $\times \left\{(23)\,(465)\,\epsilon_\xi\right\} \left\{(23654)\,\epsilon_\xi\right\}$	1 2	$\{26,1236\}$ $\{26,2456\}$	$\{12,123,46,456\}$
(16)	$\sum_{\alpha_{13}\alpha_{23}\alpha_{36}} \sum_{\alpha_{14}\alpha_{15}\alpha_{56}} \sum_{\alpha_{\zeta\in\Gamma_\xi}} f_{16} \left\{(12)\,\epsilon_6\right\}$ $\times \left\{(126)\,\epsilon_6\right\} \left\{(123)\,[\epsilon_6\epsilon_{19}]\right\} \left\{(1236)\,[\epsilon_6\epsilon_{19}]\right\}$ $\times \left\{(123)\,(56)\,\epsilon_{19}\right\} \left\{(12365)\,\epsilon_{19}\right\}$ $\times \left\{(123)\,(465)\,\epsilon_\xi\right\} \left\{(123654)\,\epsilon_\xi\right\}$	1 2	$\{16,1236\}$ $\{16,1456\}$	$\{12,123,46,456\}$

Tab. 3: Recoupling coefficients ε_{ij}

If $\widehat{\pi} = (132)$, a 3-cycle has to be decomposed into the composition of two 2-cycles. Such a decomposition is the set of operators $\widehat{\mathcal{E}}_\pi \in \left\{(13)\,(23)\,,(12)\,(13)\,,(23)\,(12)\right\}$. The quantity π in the subscript in the notation of the operator $\widehat{\mathcal{E}}_\pi$ shows that the present set has been obtained from the operator $\widehat{\pi}$. The operators $\widehat{\mathcal{E}}_\pi$ will be called the equivalent operators (or equivalent permutations) for given $\widehat{\pi}$. Particularly, $\widehat{\mathcal{E}}_{ij} = (ij)$, i.e., for a 2-cycle (ij) there is only one equivalent operator $\widehat{\mathcal{E}}_{ij}$ which coincides with (ij). Hence, $\widehat{\mathcal{E}}_{132}\left\langle iijlpq\right\rangle = \left\langle jiilpq\right\rangle = \sum_{\varsigma_{ii}=\mathrm{odd}} \varepsilon_{132}\left\langle iijlpq\right\rangle$, where $\widehat{\mathcal{E}}_{132} = \left(\begin{smallmatrix}1&2&3\\3&1&2\end{smallmatrix}\right)$. The task is to find out ε_{132}. If

$\widehat{\mathcal{E}}_{132} = (13)\,(23)$, the third operator in $\langle iijlpq\rangle$ is permuted with the second one and after that the permuted third operator is replaced with the first one (none of the permutations of the same rank operators are performed). According to Eq. (4.1), $(13)\,(23)\,\langle iijlpq\rangle$ equals to $\sum_{\varsigma_{ii}=\mathrm{odd}} \varepsilon_{13}\left\{(13)\,\varepsilon_{23}\right\}\langle iijlpq\rangle$, where $\varepsilon_{23} = \varpi_{23}\epsilon_6\left\{(23)\,\epsilon_6\right\}$. Besides, in this case identities $\varpi_{13} = \varpi_{23} = 1$. Then $(13)\,\varepsilon_{23} = \left\{(13)\,\epsilon_6\right\}\left\{(132)\,\epsilon_6\right\}$. But $\alpha_1 = \alpha_2 = \varsigma_i$, $\alpha_3 = \varsigma_j$. Then, according to Eq. (A.11) we obtain $(13)\,\varepsilon_{23} = -1$, and the coefficient $\varepsilon_{132} = -\varepsilon_{13}$. If $\widehat{\mathcal{E}}_{132} = (12)\,(13)$, then, in accordance with Eqs. (4.1), (4.3), $(12)\,(13)\,\langle iijlpq\rangle = \langle jiilpq\rangle = \sum_{\varsigma_{ii}=\mathrm{odd}}\left\{(12)\,\varepsilon_{13}\right\}\langle iijlpq\rangle = \sum_{\varsigma_{ii}=\mathrm{odd}}\varepsilon_{23}\,\langle iijlpq\rangle$. The last equality has been obtained exploiting Eq. (4.3), which says that $(12)\,\varepsilon_{13} = f_{13}f_{23}\varepsilon_{23}$. But $f_{13} = 1$, $f_{23} = \varpi_{23} = \varpi_{13} = 1$, and $(12)\,\varepsilon_{13} = \varepsilon_{23}$. On the other hand, $\varepsilon_{23} = \varepsilon_{13}\left\{(13)\,\varepsilon_{23}\right\} = -\varepsilon_{13}$. Thus, the coefficient $\varepsilon_{132} = -\varepsilon_{13}$. Finally, if $\widehat{\mathcal{E}}_{132} = (23)\,(12)$, then $(23)\,(12)\,\langle iijlpq\rangle = \sum_{\varsigma_{ii}=\mathrm{odd}}\varepsilon_{23}\,\langle iijlpq\rangle$. Hence, according to the results of the previous composition, we get that $\varepsilon_{132} = -\varepsilon_{13}$. This indicates identity $(132)\,\langle iijlpq\rangle = -\,(13)\,\langle iijlpq\rangle$. Note, (132) does not permute same rank operators, while (13) does. This means every n-length permutation $\widehat{\pi}$ with $n > 2$ can be replaced by the shorter permutation $\widehat{\pi}_{\min}$ (if such exists), i.e., the operation with the less number of the permutations, in the following way

$$\widehat{\pi}\widehat{\mathrm{T}}^{\lambda}_{\varkappa} = \theta\widehat{\pi}_{\min}\widehat{\mathrm{T}}^{\lambda}_{\varkappa}, \tag{4.6}$$

where $\theta = \pm 1$, and the sign depends whether the number of permuted same rank operators is even or odd. For example, $(1432)\,\langle iiijpq\rangle = (14)\,\langle iiijpq\rangle$, where the operation (1432) does not permute the same rank ς_i operator. The operation (14) permutes the first operator with the second and the third one. In this case $\theta = 1$. Analogous expression can be obtained if we write $(243)\,\langle iiijpq\rangle = -\,(24)\,\langle iiijpq\rangle$. Since $(14)\,(243) = (1432)$ and $(14)\,(24) = (142)$, it follows that $(1432)\,\langle iiijpq\rangle = -\,(142)\,\langle iiijpq\rangle$ is valid. On the other hand $(142)\,\langle iiijpq\rangle = -\,(14)\,\langle iiijpq\rangle$. Thus, we obtain identical result. Note, the procedures studied in this example are valid for all $\widehat{\mathcal{E}}_{1432}$. The equivalent operators $\widehat{\mathcal{E}}_{\pi}$ for $\widehat{\pi} = (ijk)$ and $\widehat{\pi} = (ijkl)$ are these

$$\begin{aligned} &\widehat{\mathcal{E}}_{ijk} \in \left\{(ij)\,(kj)\,,(kj)\,(ik)\,,(ik)\,(ij)\right\},\\ &\widehat{\mathcal{E}}_{ijkl} \in \left\{(ij)\,(jk)\,(kl)\,,(jk)\,(il)\,(ij)\,,(jk)\,(kl)\,(il)\,,(kl)\,(il)\,(ij)\right\}. \end{aligned} \tag{4.7}$$

The presented results, generalized in Eq. (4.6), demonstrate how any n-length permutation can be replaced by simpler permutation even if it permutes same rank operators. Equivalently, this fact allows one to reduce the number of summation parameters, which appear due to performed permutations.

5 Inclusion of electron-electron interaction in $n \leq 6$ open shells

In this section we address to the study of the three-particle operator $\widehat{T}^{\varsigma}_{\iota}\left(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q\right) = \langle ijklpq\rangle$ (see Eq. (2.5)), depending on how many shells $n_x\lambda_x^{N_x}$ it acts on. We distinguish operators $\langle ijklpq\rangle$ by the classes $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$, where $\Delta_x = N_x - \bar{N}_x$ denotes the difference of electron numbers in the shells $n_x\lambda_x^{N_x}$ and $n_x\lambda_x^{\bar{N}_x}$ of bra $\langle\Psi_f^x|$ and ket $|\Psi_i^x\rangle$ wave functions, respectively. It is assumed that the conservation law for electron number is satisfied

$$\sum_{x=1}^{n}\Delta_x = 0. \tag{5.1}$$

In Eq. (5.1) the summation is performed over all shells of bra and ket wave functions. The operators $\langle ijklpq\rangle$ belong to one class if the sum of numbers $x \in \{i, j, k\}$ and $\bar{x} \in \{-l, -p, -q\}$, where $x = |\bar{x}|$ (same shell), coequals to Δ_x. The dimension of class $d = \dim X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ is characterized by the number of operators $\langle ijklpq\rangle$, which belong to given class. We consider three types of classes: (i) parent; (ii) dual; (iii) derived. Each parent class $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ has its dual class $X_n^*\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$, if

$$X_n^*\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right) = X_n\left(-\Delta_1, -\Delta_2, \ldots, -\Delta_n\right), \tag{5.2}$$

$$\dim X_n^*\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right) = \dim X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right). \tag{5.3}$$

The selection which class is parent and which one is dual, is optional. It is evident, matrix elements of operators, which belong to given class and its dual one, coincide. Dual class is a particular case of derived class. Let us study derived classes in a more detail.

Suppose our task is to transform given operator $\langle ijklpq\rangle \equiv \langle x_\pi\rangle$ of $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ into a consequent order operator $\langle i'j'k'l'p'q'\rangle \equiv \langle x\rangle$, where $i' \le j' \le k' \le l' \le p' \le q'$. This transformation is depicted by permutation operator $\widehat{\pi}$ of S_n

$$\widehat{\pi}\,\langle x\rangle = \langle x_\pi\rangle. \tag{5.4}$$

Further, we demonstrate how one can obtain acquainted permutations $\widehat{\pi}'$ for $\widehat{\pi}'\langle y\rangle = \langle y_{\pi'}\rangle$, where $\langle y_{\pi'}\rangle$ belongs to $X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right)$, if the permutations $\widehat{\pi}$ of $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ are known. Suppose there is a map

$$\phi: \quad X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right) \mapsto X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right), \tag{5.5}$$

which is associated to the numbers x_π and $y_{\pi'}$ as follows

$$y_{\pi'} = \phi\left(x_\pi\right) \Rightarrow \langle y_{\pi'}\rangle = \phi\,\langle x_\pi\rangle. \tag{5.6}$$

Substituting Eq. (5.4) in Eq. (5.6) one gets

$$\widehat{\pi}\,\langle x\rangle = \phi^{-1}\widehat{\pi}'\,\langle y\rangle. \tag{5.7}$$

The last equation means that for every operation $\widehat{\pi}$ on the operator $\langle x\rangle$ from the class $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ there exists the operation $\phi^{-1}\widehat{\pi}'$ on the operator $\langle y\rangle$ from the class $X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right)$ such that the resultant operator $\langle x_\pi\rangle$ is invariant. Further, let $\widehat{\Phi}$ be defined by

$$\langle x\rangle = \widehat{\Phi}\,\langle y\rangle. \tag{5.8}$$

Then it follows from Eqs. (5.7), (5.8)

$$\widehat{\pi}' = \phi\widehat{\pi}\widehat{\Phi}. \tag{5.9}$$

Obtained expression connects the known permutation $\widehat{\pi}$ of the class $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ with the permutation $\widehat{\pi}'$ of $X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right)$, called derived class. It is important to mention, that the number of operators $\widehat{\Phi}$ is much more less than the number $\dim X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right)$ of operators $\widehat{\pi}'$. For example, $\dim X_3\left(+2, -1, -1\right) = 24$, while the number of operators $\widehat{\Phi}$ equals to 3. This number coincides with amount of the ordered operators $\langle x\rangle$ (it directly follows from Eq. (5.8)).

At this step we pick out special case $\phi = \Pi_{\mu\nu}$, where

$$\Pi_{\mu\nu}X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_\mu, \ldots, \Delta_\nu, \ldots, \Delta_n\right) = X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_\nu, \ldots, \Delta_\mu, \ldots, \Delta_n\right). \tag{5.10}$$

Then

$$\widehat{\Phi} = \Pi_{\mu\nu}\widehat{\widehat{\pi}}. \tag{5.11}$$

For instance, if $X_n\left(\Delta_1', \Delta_2', \Delta_3, \ldots, \Delta_n'\right) = X_n\left(\Delta_2, \Delta_1, \Delta_3, \ldots, \Delta_n\right)$, then $\Pi_{\mu\nu} = \Pi_{12}$. Further, if $X_n\left(\Delta_1, \Delta_2, \ldots,\right.$ $X_3\left(+2, -1, -1\right)$, then $X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right) = X_3\left(-1, +2, -1\right)$. Notice, the signs of Δ_1 and Δ_2 have changed. For this reason the operator $\Pi_{\mu\nu}\widehat{\pi}\widehat{\Phi}$ has to be replaced by $\Pi_{\mu\nu}\widehat{\Phi}\widehat{\pi}$. Then from Eqs. (5.9)-(5.11) follows

$$\widehat{\pi}' = \widehat{\widehat{\pi}}\widehat{\pi} \tag{5.12}$$

and the operator $\hat{\hat{\pi}}$ is generated from the next equation

$$\hat{\hat{\pi}} \left\langle y \right\rangle = \Pi_{\mu\nu} \left\langle x \right\rangle , \tag{5.13}$$

where $\Pi_{\mu\nu}$ permutes the numbers μ and ν in x. In Eq. (5.13) $\Pi_{\mu\nu}$ can be substituted by various products of $\Pi_{\mu_1\nu_1}\Pi_{\mu_2\nu_2}\ldots$ depending on the classes in which the operators $\langle x_\pi \rangle$ and $\langle y_{\pi'} \rangle$ are prescribed. Notice, Eq. (5.12) allows one to compute, particularly, the permutations of dual class. If, e.g., $\langle x \rangle = \langle 111112 \rangle$, then $\langle y \rangle = \langle 122222 \rangle$ and $\hat{\hat{\pi}} = (16)$. The representation of this permutation in dual class is namely $(16)\,1_6 = (16)$. Suppose $\hat{\pi} = -\,(13)$ and $\langle x \rangle = \langle 112223 \rangle$. Then $\langle x_\pi \rangle = \langle 211223 \rangle$. For $\Pi_{\mu\nu} = \Pi_{12}$, $\langle y_{\pi'} \rangle = \Pi_{12} \langle x_\pi \rangle = \langle 122113 \rangle$ and $\langle y \rangle = \langle 111223 \rangle$, where the operators $\langle x \rangle , \langle x_\pi \rangle \in X_3\left(+2,-1,-1\right)$ and $\langle y \rangle , \langle y_{\pi'} \rangle \in X_3\left(-1,+2,-1\right)$. It follows from Eq. (5.13), $\hat{\hat{\pi}} \langle 111223 \rangle = \Pi_{12} \langle 112223 \rangle = \langle 221113 \rangle \Rightarrow \hat{\hat{\pi}} = (14253)$. According to Eq. (4.6), the operator $\hat{\hat{\pi}}$ can be replaced by the less number permutation $\hat{\hat{\pi}} = \hat{\hat{\pi}}_{\min} = (14)\,(25)$. Then $\hat{\pi}' = \hat{\hat{\pi}}\hat{\pi} = -\,(14)\,(25)\,(13) = -\,(134)\,(25) = (25)\,(34)$, where the last equality has been obtained exploiting Eq. (4.6). The result is checked by the equation $\hat{\pi}' \langle y \rangle = \langle y_{\pi'} \rangle$, i.e., $\hat{\pi}' \langle 111223 \rangle = \langle 122113 \rangle \Rightarrow \hat{\pi}' = (24)\,(35) = (25)\,(34)$. Hence, both methods produce identical results. Notice, $\hat{\hat{\pi}}$ is the same for all operators $\langle x_\pi \rangle$, if the ordered operator is $\langle x \rangle$. It directly follows from Eq. (5.13). That means it is enough to find out operators $\hat{\hat{\pi}}$ - all other permutations $\hat{\pi}'$ of derived class $X_n\left(\Delta_1', \Delta_2', \ldots, \Delta_n'\right)$ are found from Eq. (5.12), if the permutations $\hat{\pi}$ of parent class $X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$ are known.

The classification of irreducible tensor operators $\langle ijklpq \rangle$ is presented in Appendix B in Tabs. 8-23. In tables notation $\langle ijk\left\{lpq\right\} \rangle$ denotes the set $\left\{\langle ijklpq \rangle , \langle ijklqp \rangle , \langle ijkplq \rangle , \langle ijkpql \rangle , \langle ijkqlp \rangle , \langle ijkqpl \rangle\right\}$. Every permutation of the numbers in the set $\{lpq\}$ is prescribed by the operation $\hat{\vartheta}$, where

$$\hat{\vartheta} \in \left\{1_6, (56)\,, (45)\,, (456)\,, (465)\,, (46)\right\} . \tag{5.14}$$

Similarly, $\langle \{ijk\}\,lpq \rangle$ denotes the set $\left\{\langle ijklpq \rangle , \langle ikjlpq \rangle , \langle jiklpq \rangle , \langle jkilpq \rangle , \langle kijlpq \rangle , \langle kjilpq \rangle\right\}$, and every permutation of the numbers in the set $\{ijk\}$ is prescribed by the operation $\hat{\eta}$, where

$$\hat{\eta} \in \left\{1_6, (23)\,, (12)\,, (123)\,, (132)\,, (13)\right\} . \tag{5.15}$$

The operators $\hat{\eta}$ and $\hat{\vartheta}$ commute with each other, i.e., $[\hat{\eta}, \hat{\vartheta}] = 0$.

One fact needs to be mentioned. If some of the differences Δ_x and Δ_y coequal, the expressions of $\hat{\hat{\pi}}$ can acquire different forms, though they are equivalent. Suppose (see Tab. 20), that $\langle y \rangle = \langle 122344 \rangle \in X_4\left(\Delta_1', \Delta_2', \Delta_3', \Delta_4'\right) = X_4\left(-1, 0, -1, +2\right)$. Then operator $\hat{\hat{\pi}} = -\,(1634)\,(25)$ and it satisfies Eq. (5.13) with $\Pi_{\mu\nu} = \Pi_{23}\Pi_{14}\Pi_{34}$. Further, if $\langle x_\pi \rangle = \langle 114234 \rangle$, then $\hat{\pi} = (354)\,1_6 = (354)$. Consequently, permutation $\hat{\pi}' = \hat{\hat{\pi}}\hat{\pi} = -\,(1634)\,(25)\,(354) = -\,(15)\,(26)$ and thus irreducible tensor operator equals to $\langle y_{\pi'} \rangle = \langle 442132 \rangle \in X_4\left(-1, 0, -1, +2\right)$. In this case $\Delta_2 = \Delta_3 = -1$ and $\Delta_1' = \Delta_3' = -1$. Hence there is another $\hat{\hat{\pi}} = -\,(163)\,(25)$ value, which also satisfies Eq. (5.13) with $\Pi_{\mu\nu} = \Pi_{14}\Pi_{12}\Pi_{24}$. This implies $\hat{\pi}' = \hat{\hat{\pi}}\hat{\pi} = -\,(163)\,(25)\,(354) = -\,(154)\,(26)$ and the tensor operator $\langle y_{\pi'} \rangle = \langle 442312 \rangle \in X_4\left(-1, 0, -1, +2\right)$. It is seen, both operators $\langle 442132 \rangle$ and $\langle 442312 \rangle$ belong to the same class $X_4\left(-1, 0, -1, +2\right)$, where $\langle 442132 \rangle = (45)\,\langle 442312 \rangle$. Hence, both $\hat{\hat{\pi}}$ operators generate the elements of same class.

6 Matrix elements

In this section we study matrix elements $\langle \Psi_f |\, {}_{\Lambda}\hat{T}^{\lambda}_{\mu} | \Psi_i \rangle$ on the basis[3]

$$|\Psi_i\rangle = \left| \lambda_1^{N_1} \lambda_2^{N_2} \ldots \lambda_n^{N_n} \Gamma_1\Lambda_1\Gamma_2\Lambda_2\Lambda_{12}\Gamma_3\Lambda_3\Lambda_{123}\ldots\Gamma_u\Lambda_u\Lambda M_\Lambda \right\rangle , \tag{6.1}$$

composed of u shells of equivalent electrons, where single-shell function in Λ-space is defined by

$$|\Psi_i^x\rangle = \left| \lambda_x^{N_x} \Gamma_x \Lambda_x M_{\Lambda_x} \right\rangle , \tag{6.2}$$

---

[3] The subscripts i and f in Ψ_i and Ψ_f designate initial and final states.

with Γ_x indicating additional quantum numbers. In q-space $|\Psi_i^x\rangle$ is denoted $|\lambda_x\Gamma_x q_x m_{q_x}\rangle$ ($q \equiv Q\Lambda$). Then the total wave function $|\Psi_i\rangle$ in q-space (for $n_x = \text{const.}\ \forall x = 1, 2, \ldots, u$) is represented by superposition of functions $|\lambda_x\Gamma_x q_x m_{q_x}\rangle$, coupled by the same scheme as in Eq. (6.1). Namely,

$$|\left(\lambda_1 + \lambda_2 + \ldots + \lambda_u\right)\Gamma_1 q_1 \Gamma_2 q_2 q_{12} \ldots \Gamma_u q_u q m_q\rangle\,. \tag{6.3}$$

The momenta coupling is performed by using known rules of angular momentum theory. The coupling coefficients are usual Clebsch-Gordan coefficients. In Eq. (6.3) the representations λ_x are written in a sum differently from Eq. (6.1). This is because the total basis index of Q in Eq. (6.3) is $M_Q = M_{Q_1} + M_{Q_2} + \ldots + M_{Q_u} = \left(2N - [\lambda]\right)/4$, where $N = N_1 + N_2 + \ldots + N_u$ and $\lambda = \lambda_1 + \lambda_2 + \ldots + \lambda_u$, i.e., the representation of constructed function in Λ-space is the function of the shell $n_x\lambda^N$ [9].

The expressions of reduced matrix elements in quasispin formalism can be found in various literature. See, for instance, [7, 10]. Reduced matrix element of operator ${}_{\Lambda}\widehat{T}^{\lambda}$ in Λ-space will be marked by $\left[\lambda_x^{N_x}\Gamma_x\Lambda_x\|\,{}_{\Lambda}\widehat{T}^{\lambda}\|\lambda_x^{\bar{N}_x}\bar{\Gamma}_x\bar{\Lambda}_x\right]$, while reduced matrix element of operator $\widehat{T}^{\varsigma}$ in q-space will be denoted by $\left[\lambda_x\Gamma_x q_x|||\widehat{T}^{\varsigma}|||\lambda_x\bar{\Gamma}_x\bar{q}_x\right]$.

The relationship between reduced matrix elements in Λ and q spaces of the corresponding operators ${}_{\Lambda}\widehat{T}^{\lambda}$ and $\widehat{T}^{\varsigma} = \widehat{T}^{\kappa\lambda}$ (see Eqs. (2.4)-(2.5)) is associated by the next equation

$$\begin{aligned}&\left[\lambda_1^{N_1}\lambda_2^{N_2}\ldots\lambda_u^{N_u}\Gamma_1\Lambda_1\Gamma_2\Lambda_2\Lambda_{12}\ldots\Gamma_u\Lambda_u\Lambda\|\,{}_{\Lambda}\widehat{T}^{\lambda}\|\lambda_1^{\bar{N}_1}\lambda_2^{\bar{N}_2}\ldots\lambda_u^{\bar{N}_u}\bar{\Gamma}_1\bar{\Lambda}_1\bar{\Gamma}_2\bar{\Lambda}_2\bar{\Lambda}_{12}\ldots\bar{\Gamma}_u\bar{\Lambda}_u\bar{\Lambda}\right]\\ &= 6\delta_{N,\bar{N}}\delta_{\pi,0}\delta_{\kappa_{ij},1}\delta_{\kappa_{ij},\kappa_{lp}}\delta_{\kappa_{ijk},\frac{3}{2}}\delta_{\kappa_{ijk},\kappa_{lpq}}\sum_{\kappa=0}^{3}\frac{[\kappa]^{\frac{1}{2}}}{\sqrt{(3-\kappa)!\,(4+\kappa)!}}\left\langle\bar{Q}M_Q\kappa 0|QM_Q\right\rangle\\ &\times\left[\left(\lambda_1+\lambda_2+\ldots+\lambda_u\right)\Gamma_1 q_1\Gamma_2 q_2 q_{12}\ldots\Gamma_u q_u q|||\widehat{T}^{\kappa\lambda}|||\left(\lambda_1+\lambda_2+\ldots+\lambda_u\right)\bar{\Gamma}_1\bar{q}_1\bar{\Gamma}_2\bar{q}_2\bar{q}_{12}\ldots\bar{\Gamma}_u\bar{q}_u\bar{q}\right].\end{aligned} \tag{6.4}$$

Note, the condition $\pi = 0$ has already been used in Sec. 5, when classifying tensor operators $\langle ijklpq\rangle$. It directly follows from Eq. (6.4) that $N = \bar{N}$, where $N = \sum_{x=1}^{u} N_x$ and $\bar{N} = \sum_{x=1}^{u}\bar{N}_x$. This coincides with Eq. (5.1).

Reduced matrix element of $\widehat{T}^{\varsigma}\left(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q\right)$ with $i, j, k, l, p, q \in \{1, 2, \ldots, n\} \subseteq \mathcal{I}$ when $n > 1$ is constructed as follows

$$\begin{aligned}&\left[\left(\lambda_1+\ldots+\lambda_n\right)\Gamma_1 q_1\Gamma_2 q_2 q_{12}\ldots\Gamma_n q_n q|||\widehat{T}^{\varsigma}\left(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q\right)|||\left(\lambda_1+\ldots+\lambda_n\right)\bar{\Gamma}_1\bar{q}_1\bar{\Gamma}_2\bar{q}_2\bar{q}_{12}\ldots\bar{\Gamma}_n\bar{q}_n\bar{q}\right]\\ &= (-1)^{\Phi_n}\sum_{\varsigma_w\in\mathcal{L}_\xi}\epsilon_\xi\prod_{x=1}^{n}\left[q_{12\ldots x}, q_{x+1}, \bar{q}_{12\ldots x+1}, \varsigma_{p_{x+1}}\right]^{\frac{1}{2}}\begin{Bmatrix}\bar{q}_{12\ldots x} & \bar{q}_{x+1} & \bar{q}_{12\ldots x+1}\\ \varsigma_{p_x} & \varsigma_{p_{x+1\,x+1}} & \varsigma_{p_{x+1}}\\ q_{12\ldots x} & q_{x+1} & q_{12\ldots x+1}\end{Bmatrix}\\ &\times\left[\lambda_x\Gamma_x q_x|||\widehat{\mathcal{O}}^{\varsigma_{p_{xx}}}\left(\lambda_x\lambda_x\ldots\lambda_x\right)|||\lambda_x\bar{\Gamma}_x\bar{q}_x\right];\quad q_{12\ldots n} = q,\quad \bar{q}_{12\ldots n} = \bar{q},\quad \varsigma_{p_n} = \varsigma.\end{aligned} \tag{6.5}$$

The numbers p_x and $p_{\ell\ell}$ are defined by

$$p_x = 11\ldots 122\ldots 2xx\ldots x,\quad p_{\ell\ell} = \ell\ell\ldots\ell\quad (\mathcal{N}_\ell\,\text{times})\,. \tag{6.6}$$

The phase multiplier Φ_n is

$$\Phi_n = \sum_{x=1}^{n}\left[\left(N_x + \mathcal{N}_x + \Delta_x\right)\sum_{y>x}^{n} N_y + \mathcal{N}_x\sum_{z=1}^{x-1}\left(N_z + \Delta_z\right)\right]. \tag{6.7}$$

Here it is assumed that $\widehat{T}^{\varsigma}\left(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q\right)$ is presented in a consequent order (Sec. 5), i.e., it is of the form $\langle x\rangle \in X_n\left(\Delta_1, \Delta_2, \ldots, \Delta_n\right)$. In Eq. (6.7) $\mathcal{N}_x$ denotes the number of operators a^{ς_x} in $\widehat{T}^{\varsigma}\left(\lambda_i\lambda_j\lambda_k\lambda_l\lambda_p\lambda_q\right)$. In general, the operator $\widehat{\mathcal{O}}$ reads

$X_2\,(\Delta_1,\Delta_2)$	$\langle x\rangle$	ξ	w_1	w_2
$X_2\,(0,0)$	$\langle 111122\rangle$	20	1111	22
	$\langle 112222\rangle$	5	2222	–
$X_2\,(+1,-1)$	$\langle 111112\rangle$	27	1111	11111
	$\langle 111222\rangle$	14	–	–
	$\langle 122222\rangle$	30	2222	22222

Tab. 4: The parameters for three-particle matrix element. Two-shell case

$X_3\,(\Delta_1,\Delta_2,\Delta_3)$	$\langle x\rangle$	ξ	w_1	w_2	w_3
$X_3\,(0,0,0)$	$\langle 112233\rangle$	1	22	33	1122
$X_3\,(+2,-1,-1)$	$\langle 111123\rangle$	27	1111	11112	–
	$\langle 112223\rangle$	4	222	11222	–
	$\langle 112333\rangle$	14	–	–	–
$X_3\,(+3,-2,-1)$	$\langle 111223\rangle$	15	11122	–	–
$X_3\,(+1,-1,0)$	$\langle 123333\rangle$	5	3333	–	–
	$\langle 111233\rangle$	20	1112	33	–
	$\langle 122233\rangle$	24	22	33	222
$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$	$\langle y\rangle$	ξ	w_1	w_2	w_3
$X_3\,(-1,+2,-1)$	$\langle 122223\rangle$	31	222	2222	12222
	$\langle 122333\rangle$	6	22	–	–

Tab. 5: The parameters for three-particle matrix element. Three-shell case

$$\widehat{\mathcal{O}}^{\varsigma_{xy\ldots z}}\left(\lambda_x\lambda_y\ldots\lambda_z\right) = [[\ldots[a^{\varsigma_x}\times a^{\varsigma_y}]^{\varsigma_{xy}}\times\ldots]^{\varsigma_{xy\ldots}}\times a^{\varsigma_z}]^{\varsigma_{xy\ldots z}}\,. \tag{6.8}$$

The representation of $\widehat{\mathcal{O}}^{\varsigma}$ in Λ-space is $\widehat{\mathcal{O}}^{\lambda}$. Reduced matrix elements of $\widehat{\mathcal{O}}^{\lambda}$ are known. Expressions can be found, for instance, in [11]. Thus, the reduced matrix element on the right hand side of Eq. (6.5) is assumed to be known. In Eq. (6.5) operators $\widehat{\mathcal{O}}^{\varsigma_{p_{xx}}}\left(\lambda_x\lambda_x\ldots\lambda_x\right)$ consist of $\mathcal{N}_x$ operators a^{ς_x}. Representations $\varsigma_{p_{\ell\ell}}$ with $p_{\ell\ell}$ being $\mathcal{N}_\ell$-length number, are prescribed in q_ℓ-space. The summation is performed over ς_w, where the indices $w\in\mathcal{L}_\xi=\{w_1,w_2,w_3\}$. The values of $w_{i\in\{1,2,3\}}$ depend on the concrete operator $\langle x\rangle$. These are depicted in Tabs. 4-7.

$X_4\,(\Delta_1,\Delta_2,\Delta_3,\Delta_4)$	$\langle x\rangle$	ξ	w_1	w_2	w_3
$X_4\,(+1,+1,-1,-1)$	$\langle 111234\rangle$	27	1112	11123	–
	$\langle 122234\rangle$	29	222	1222	12223
	$\langle 123334\rangle$	4	333	12333	–
	$\langle 123444\rangle$	14	–	–	–
$X_4\,(+2,-2,+1,-1)$	$\langle 112234\rangle$	3	22	1122	11223
$X_4\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4)$	$\langle y\rangle$	ξ	w_1	w_2	w_3
$X_4\,(+2,+1,-2,-1)$	$\langle 112334\rangle$	15	11233	–	–
$X_4\,(+2,+1,-1,-2)$	$\langle 112344\rangle$	20	44	1123	–
$X_4\,(+1,+2,-2,-1)$	$\langle 122334\rangle$	7	12233	–	–
$X_4\,(+1,+2,-1,-2)$	$\langle 122344\rangle$	23	44	1223	–
$X_4\,(+1,-1,+2,-2)$	$\langle 123344\rangle$	1	33	44	1233

Tab. 6: The parameters for three-particle matrix element. Four-shell case

$X_5\,(\Delta_1,\Delta_2,\Delta_3,\Delta_4,\Delta_5)$	$\langle x\rangle$	ξ	w_1	w_2	w_3
$X_5\,(+2,+1,-1,-1,-1)$	$\langle 112345\rangle$	27	1123	11234	–
$X_5\,(+1,+1,-1,-1,0)$	$\langle 123455\rangle$	20	55	1234	–
$X_5\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4,\Delta'_5)$	$\langle y\rangle$	ξ	w_1	w_2	w_3
$X_5\,(+1,+2,-1,-1,-1)$	$\langle 122345\rangle$	28	22	1223	12234
$X_5\,(-1,-1,+2,+1,-1)$	$\langle 123345\rangle$	3	33	1233	12334
$X_5\,(-1,+1,-1,+2,-1)$	$\langle 123445\rangle$	15	12344	–	–

Tab. 7: The parameters for three-particle matrix element. Five-shell case

When $n=6$, the parameters, suitable for reduced matrix element of $\langle x\rangle=\langle 123456\rangle$, where $\langle 123456\rangle\in X_6\,(+1,+1,+1,-1,-1,-1)$, are these: $\xi=27$, $w_1=1234$, $w_2=12345$.

Let us take some examples, which show how one can easily obtain the structure of $\langle x\rangle$ matrix element. Suppose $\langle x\rangle = \langle 111122\rangle$. Then $\mathcal{N}_1 = 4$, $\mathcal{N}_2 = 2$ and (see Tab. 4)

$$\left[(\lambda_1+\lambda_2)\,\Gamma_1 q_1 \Gamma_2 q_2 q|||\widehat{T}^{\varsigma}\,(\lambda_1\lambda_1\lambda_1\lambda_1\lambda_2\lambda_2)\,|||\,(\lambda_1+\lambda_2)\,\bar{\Gamma}_1\bar{q}_1\bar{\Gamma}_2\bar{q}_2\bar{q}\right] = (-1)^{N_1N_2}\,[q_1,q_2,\bar{q},\varsigma]^{\frac{1}{2}}\sum_{\varsigma_{1111}\varsigma_{22}}\epsilon_{20}$$
$$\times\begin{Bmatrix}\bar{q}_1 & \bar{q}_2 & \bar{q}\\ \varsigma_{1111} & \varsigma_{22} & \varsigma\\ q_1 & q_2 & q\end{Bmatrix}\left[\lambda_1\Gamma_1 q_1|||\widehat{\mathcal{O}}^{\varsigma_{1111}}\,(\lambda_1\lambda_1\lambda_1\lambda_1)\,|||\lambda_1\bar{\Gamma}_1\bar{q}_1\right]\left[\lambda_2\Gamma_2 q_2|||\widehat{\mathcal{O}}^{\varsigma_{22}}\,(\lambda_2\lambda_2)\,|||\lambda_2\bar{\Gamma}_2\bar{q}_2\right]. \qquad (6.9)$$

Other matrix elements of $\langle x_\pi\rangle = \widehat{\pi}\,\langle x\rangle$ are obtained using Eq. (4.5). For example, the operator $\langle 121121\rangle = -(26)\,\langle 111122\rangle$. This points to (see Eq. (4.1) and Tab. 8)

$$\left[(\lambda_1+\lambda_2)\,\Gamma_1 q_1 \Gamma_2 q_2 q|||\widehat{T}^{\varsigma}\,(\lambda_1\lambda_2\lambda_1\lambda_1\lambda_2\lambda_1)\,|||\,(\lambda_1+\lambda_2)\,\bar{\Gamma}_1\bar{q}_1\bar{\Gamma}_2\bar{q}_2\bar{q}\right]$$
$$= -\sum_{\varsigma_{11}\varsigma_{111}\varsigma_{122}}\varepsilon_{26}\left[(\lambda_1+\lambda_2)\,\Gamma_1 q_1 \Gamma_2 q_2 q|||\widehat{T}^{\varsigma}\,(\lambda_1\lambda_1\lambda_1\lambda_1\lambda_2\lambda_2)\,|||\,(\lambda_1+\lambda_2)\,\bar{\Gamma}_1\bar{q}_1\bar{\Gamma}_2\bar{q}_2\bar{q}\right]. \qquad (6.10)$$

7 Conclusions

The comprehensive treatment of an effective three-particle operator $\widehat{L}_3$ for open-shell atoms was presented. However, for open-shell atoms the overall treatment of this operator is very complex and tedious. In present paper effective three-particle operator was studied requesting quasispin formalism. Due to the complexity and high abundance of miscellaneous forms to reduce given operator $\widehat{L}_3$, the procedure, suitable for any $\widehat{L}_n$, to classify irreducible tensor operators were performed by exploiting irreducible representations of S_n, combined with proposed adaptation of n-tuple concept. Recoupling coefficients, called the basis coefficients, which arise when transforming one irreducible tensor form into another, are expressed by $3nj$-coefficients (Appendix A). As long as $\widehat{L}_3$ consists of six operators a^{ς_x}, there are $46,656$ ways to express chosen coupling scheme with respect to the ordering of ranks ς_x. These possible distributions were also determined proposing the so-called permutation coefficients, i.e., the coefficients which arise when transforming one partition with its own ordering of ranks into another (or the same) partition with different ordering of ranks. Permutation coefficients, expressed by the basis coefficients, appear when a 2-cycle operation (ij) acts on chosen tensor operator in such a way, that the ordering of given operator ranks is permuted into the consequent order, labeled by the sequence $1,2,\ldots,6$. The permutation considered makes sense when one-shell wave functions, on the basis of which the matrix element is calculated, are coupled in analogous consequent order. Permutation coefficients were constructed making the least number of recouplings, since, in general, every permutation stipulates the summation over intermediate ranks. Besides, we have proved that every permutation can be replaced by the less number permutation (if such exists), irrespective of whether the ranks of permuted operators a^{ς_x} are equal or not. Present circumstance allows to reduce the amount of summation parameters, which are found when performing recoupling of ranks.

The three-particle operators, which act on $n = 1,2,\ldots,6$ shells of equivalent electrons, were distinguished by classes (see Sec. 5 and Appendix B). These classes are characterized by the number of electrons in bra and ket wave functions, and by the number of shells. The proposed classification allows us to find out momenta recoupling coefficients, which appear when permuting the ranks of any tensor operator from total number $46,656$. Consequently, proposed systematic classification permits to account for electron-electron correlations in n open shells ($n \leq 6$) in a convenient way. The inclusion of these effects is estimated by three-particle matrix elements (Sec. 6).

Appendices

A Basis coefficients

$$\begin{aligned}\epsilon_1 &= (-1)^{\wp_1}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\left[\alpha_{34},\alpha_{45},\alpha_{56},\alpha_{123},\alpha_{456},\alpha_{1234}\right]^{\frac{1}{2}} \\ &\times \left\{\begin{matrix}\alpha_6 & \alpha_5 & \alpha_{56} \\ \alpha_4 & \alpha_{456} & \alpha_{45}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_4 & \alpha_{1234} & \alpha_{34}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{123} & \alpha_4 & \alpha_{1234} \\ \alpha_{56} & \alpha & \alpha_{456}\end{matrix}\right\},\end{aligned} \tag{A.1}$$

$$\wp_1 = \alpha_5+\alpha_6-\alpha_{56}+\alpha-\alpha_{12}+\alpha_{123}-\alpha_{1234}-\alpha_3-\alpha_4-\alpha_{456}. \tag{A.2}$$

$$\begin{aligned}\epsilon_2 &= (-1)^{\wp_2}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\left[\alpha_{34},\alpha_{45},\alpha_{56},\alpha_{123},\alpha_{456},\alpha_{3456}\right]^{\frac{1}{2}} \\ &\times \left\{\begin{matrix}\alpha_6 & \alpha_5 & \alpha_{56} \\ \alpha_4 & \alpha_{456} & \alpha_{45}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456}\end{matrix}\right\}\left\{\begin{matrix}\alpha_3 & \alpha_4 & \alpha_{34} \\ \alpha_{56} & \alpha_{3456} & \alpha_{456}\end{matrix}\right\},\end{aligned} \tag{A.3}$$

$$\wp_2 = \alpha_5+\alpha_6-\alpha_{56}+\alpha+\alpha_{12}+2\alpha_3+\alpha_{3456}. \tag{A.4}$$

$$\begin{aligned}\epsilon_3 &= (-1)^{\wp_3}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\left[\alpha_{34},\alpha_{45},\alpha_{123},\alpha_{456},\alpha_{1234},\alpha_{12345}\right]^{\frac{1}{2}} \\ &\times \left\{\begin{matrix}\alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_4 & \alpha_{1234} & \alpha_{34}\end{matrix}\right\}\left\{\begin{matrix}\alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{12345} & \alpha_{123} & \alpha_{1234}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345}\end{matrix}\right\},\end{aligned} \tag{A.5}$$

$$\wp_3 = -\alpha_{45}+\alpha_{12345}-\alpha-\alpha_5+\alpha_6-\alpha_{12}-\alpha_{1234}-\alpha_3. \tag{A.6}$$

$$\begin{aligned}\epsilon_4 &= (-1)^{\wp_4}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\left[\alpha_{34},\alpha_{45},\alpha_{123},\alpha_{345},\alpha_{456},\alpha_{12345}\right]^{\frac{1}{2}} \\ &\times \left\{\begin{matrix}\alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{45} & \alpha_{12345} & \alpha_{345}\end{matrix}\right\}\left\{\begin{matrix}\alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{345} & \alpha_3 & \alpha_{34}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345}\end{matrix}\right\},\end{aligned} \tag{A.7}$$

$$\wp_4 = -\alpha_4-\alpha_5+\alpha_{345}-\alpha-\alpha_{12}-\alpha_{123}-\alpha_{12345}-\alpha_6. \tag{A.8}$$

$$\begin{aligned}\epsilon_5 &= (-1)^{\wp_5}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\left[\alpha_{34},\alpha_{45},\alpha_{123},\alpha_{345},\alpha_{456},\alpha_{3456}\right]^{\frac{1}{2}} \\ &\times \left\{\begin{matrix}\alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456}\end{matrix}\right\}\left\{\begin{matrix}\alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{345} & \alpha_3 & \alpha_{34}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha_{3456} & \alpha_3 & \alpha_{345}\end{matrix}\right\},\end{aligned} \tag{A.9}$$

$$\wp_5 = -\alpha_3-\alpha_4+\alpha_5+\alpha_6+\alpha_{45}-\alpha_{12}+\alpha_{345}-\alpha_{456}-\alpha_{3456}-\alpha. \tag{A.10}$$

$$\begin{aligned}\epsilon_6 &= (-1)^{\wp_6}\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)\Delta\left(\alpha_{45},\alpha_6,\alpha_{456}\right)\Delta\left(\alpha_{123},\alpha_{456},\alpha\right) \\ &\times \left[\alpha_{12},\alpha_{23}\right]^{\frac{1}{2}}\left\{\begin{matrix}\alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23}\end{matrix}\right\},\end{aligned} \tag{A.11}$$

$$\wp_6 = \alpha_1+\alpha_2+\alpha_3+\alpha_{123}. \tag{A.12}$$

$$\begin{aligned}\epsilon_7 &= (-1)^{\wp_7}\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)\left[\alpha_{12},\alpha_{23},\alpha_{456},\alpha_{12345}\right]^{\frac{1}{2}} \\ &\times \left\{\begin{matrix}\alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23}\end{matrix}\right\}\left\{\begin{matrix}\alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345}\end{matrix}\right\},\end{aligned} \tag{A.13}$$

$$\wp_7 = \alpha_1+\alpha_2+\alpha_3-\alpha_6-\alpha_{45}-\alpha. \tag{A.14}$$

$$\epsilon_8 = (-1)^{\wp_8}\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)\left[\alpha_{12},\alpha_{23},\alpha_{123},\alpha_{456},\alpha_{2345},\alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times\left\{\begin{array}{ccc}\alpha_1&\alpha_2&\alpha_{12}\\\alpha_3&\alpha_{123}&\alpha_{23}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_1&\alpha_{23}&\alpha_{123}\\\alpha_{45}&\alpha_{12345}&\alpha_{2345}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_{45}&\alpha_6&\alpha_{456}\\\alpha&\alpha_{123}&\alpha_{12345}\end{array}\right\}, \tag{A.15}$$

$$\wp_8=\alpha_2+\alpha_3-\alpha_{12345}+2\alpha_1+\alpha_{23}+\alpha+\alpha_6. \tag{A.16}$$

$$\epsilon_9=(-1)^{\wp_9}\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)\left[\alpha_{12},\alpha_{23},\alpha_{123},\alpha_{456},\alpha_{2345},\alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times\left\{\begin{array}{ccc}\alpha_1&\alpha_2&\alpha_{12}\\\alpha_3&\alpha_{123}&\alpha_{23}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_1&\alpha_{23}&\alpha_{123}\\\alpha_{456}&\alpha&\alpha_{23456}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_{45}&\alpha_6&\alpha_{456}\\\alpha_{23456}&\alpha_{23}&\alpha_{2345}\end{array}\right\}, \tag{A.17}$$

$$\wp_9=\alpha_2+\alpha_3+\alpha_{123}+\alpha_{23456}+\alpha_6+\alpha_{45}-\alpha_{456}-\alpha. \tag{A.18}$$

$$\epsilon_{10}=(-1)^{\wp_{10}}\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)\Delta\left(\alpha_{45},\alpha_6,\alpha_{456}\right)\left[\alpha_{12},\alpha_{23},\alpha_{123},\alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times\left\{\begin{array}{ccc}\alpha_1&\alpha_2&\alpha_{12}\\\alpha_3&\alpha_{123}&\alpha_{23}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_1&\alpha_{23}&\alpha_{123}\\\alpha_{456}&\alpha&\alpha_{23456}\end{array}\right\}, \tag{A.19}$$

$$\wp_{10}=\alpha-\alpha_{123}+\alpha_2-\alpha_{23}+\alpha_3+\alpha_{456}. \tag{A.20}$$

$$\epsilon_{11}=(-1)^{\wp_{11}}\left[\alpha_{12},\alpha_{34},\alpha_{45},\alpha_{56},\alpha_{123},\alpha_{456},\alpha_{234},\alpha_{1234}\right]^{\frac{1}{2}}$$

$$\times\left\{\begin{array}{ccc}\alpha_1&\alpha_2&\alpha_{12}\\\alpha_{34}&\alpha_{1234}&\alpha_{234}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_{12}&\alpha_3&\alpha_{123}\\\alpha_4&\alpha_{1234}&\alpha_{34}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_4&\alpha_5&\alpha_{45}\\\alpha_6&\alpha_{456}&\alpha_{56}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_4&\alpha_{56}&\alpha_{456}\\\alpha&\alpha_{123}&\alpha_{1234}\end{array}\right\}, \tag{A.21}$$

$$\wp_{11}=-\alpha_1+\alpha_2+\alpha_3+\alpha_{12}+\alpha_{34}+\alpha_5+\alpha_6-\alpha_{56}+\alpha_{456}-\alpha_4-\alpha_{123}-\alpha. \tag{A.22}$$

$$\epsilon_{12}=(-1)^{\wp_{12}}\left[\alpha_{12},\alpha_{34},\alpha_{45},\alpha_{56},\alpha_{123},\alpha_{234},\alpha_{456},\alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times\left\{\begin{array}{ccc}\alpha_4&\alpha_5&\alpha_{45}\\\alpha_6&\alpha_{456}&\alpha_{56}\end{array}\right\}\left[\begin{array}{cccc}\alpha_2&\alpha_{23456}&\alpha_3&\alpha_{456}\\\alpha_{12}&\alpha_{34}&\alpha&\alpha_{56}\\\alpha_1&\alpha_{123}&\alpha_{234}&\alpha_4\end{array}\right], \tag{A.23}$$

$$\wp_{12}=\alpha_5+\alpha_6-\alpha_{123}-\alpha_4-\alpha_{234}+\alpha_{23456}+\alpha_{12}-\alpha_{34}. \tag{A.24}$$

$$\epsilon_{13}=(-1)^{\wp_{13}}\left[\alpha_{12},\alpha_{34},\alpha_{45},\alpha_{56},\alpha_{123},\alpha_{456},\alpha_{3456},\alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times\left\{\begin{array}{ccc}\alpha_1&\alpha_2&\alpha_{12}\\\alpha_{3456}&\alpha&\alpha_{23456}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_{12}&\alpha_3&\alpha_{123}\\\alpha_{456}&\alpha&\alpha_{3456}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_4&\alpha_5&\alpha_{45}\\\alpha_6&\alpha_{456}&\alpha_{56}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_4&\alpha_{56}&\alpha_{456}\\\alpha_{3456}&\alpha_3&\alpha_{34}\end{array}\right\}, \tag{A.25}$$

$$\wp_{13}=\alpha_5+\alpha_6-\alpha_{56}-\alpha_1-\alpha_2+\alpha_{12}+2\alpha_3. \tag{A.26}$$

$$\epsilon_{14}=\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\Delta\left(\alpha_{12},\alpha_3,\alpha_{123}\right)\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)\Delta\left(\alpha_{45},\alpha_6,\alpha_{456}\right)\Delta\left(\alpha_{123},\alpha_{456},\alpha\right). \tag{A.27}$$

$$\epsilon_{15}=(-1)^{\wp_{15}}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\Delta\left(\alpha_{12},\alpha_3,\alpha_{123}\right)\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)$$

$$\times\left[\alpha_{456},\alpha_{12345}\right]^{\frac{1}{2}}\left\{\begin{array}{ccc}\alpha_{45}&\alpha_6&\alpha_{456}\\\alpha&\alpha_{123}&\alpha_{12345}\end{array}\right\}, \tag{A.28}$$

$$\wp_{15}=\alpha+\alpha_{123}+\alpha_6+\alpha_{45}. \tag{A.29}$$

$$\epsilon_{16}=(-1)^{\wp_{16}}\Delta\left(\alpha_1,\alpha_2,\alpha_{12}\right)\Delta\left(\alpha_4,\alpha_5,\alpha_{45}\right)$$

$$\times\left[\alpha_{123},\alpha_{345},\alpha_{456},\alpha_{12345}\right]^{\frac{1}{2}}\left\{\begin{array}{ccc}\alpha_{12}&\alpha_3&\alpha_{123}\\\alpha_{45}&\alpha_{12345}&\alpha_{345}\end{array}\right\}\left\{\begin{array}{ccc}\alpha_{45}&\alpha_6&\alpha_{456}\\\alpha&\alpha_{123}&\alpha_{12345}\end{array}\right\}, \tag{A.30}$$

$$\wp_{16} = \alpha - \alpha_{12} + \alpha_{123} - \alpha_{12345} - \alpha_3 + \alpha_6. \tag{A.31}$$

$$\begin{aligned} \epsilon_{17} &= (-1)^{\wp_{17}} \Delta\left(\alpha_1, \alpha_2, \alpha_{12}\right) \Delta\left(\alpha_4, \alpha_5, \alpha_{45}\right) \\ &\times \left[\alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{3456}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix} \begin{Bmatrix} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha_{3456} & \alpha_3 & \alpha_{345} \end{Bmatrix}, \end{aligned} \tag{A.32}$$

$$\wp_{17} = \alpha + \alpha_{12} - \alpha_{3456} + \alpha_{45} - \alpha_{456} + \alpha_6. \tag{A.33}$$

$$\begin{aligned} \epsilon_{18} &= (-1)^{\wp_{18}} \Delta\left(\alpha_1, \alpha_2, \alpha_{12}\right) \Delta\left(\alpha_4, \alpha_5, \alpha_{45}\right) \Delta\left(\alpha_{45}, \alpha_6, \alpha_{456}\right) \\ &\times \left[\alpha_{123}, \alpha_{3456}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix}, \end{aligned} \tag{A.34}$$

$$\wp_{18} = \alpha + \alpha_{12} + \alpha_3 + \alpha_{456}. \tag{A.35}$$

$$\begin{aligned} \epsilon_{19} &= (-1)^{\wp_{19}} \Delta\left(\alpha_1, \alpha_2, \alpha_{12}\right) \Delta\left(\alpha_{12}, \alpha_3, \alpha_{123}\right) \Delta\left(\alpha_{123}, \alpha_{456}, \alpha\right) \\ &\times \left[\alpha_{45}, \alpha_{56}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix}, \end{aligned} \tag{A.36}$$

$$\wp_{19} = \alpha_4 + \alpha_5 + \alpha_6 + \alpha_{456}. \tag{A.37}$$

$$\begin{aligned} \epsilon_{20} &= (-1)^{\wp_{20}} \Delta\left(\alpha_1, \alpha_2, \alpha_{12}\right) \Delta\left(\alpha_{12}, \alpha_3, \alpha_{123}\right) \\ &\times \left[\alpha_{45}, \alpha_{56}, \alpha_{456}, \alpha_{1234}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_{56} & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{1234} \end{Bmatrix}, \end{aligned} \tag{A.38}$$

$$\wp_{20} = \alpha + \alpha_{123} - \alpha_{456} + \alpha_5 + \alpha_6 - \alpha_{56}. \tag{A.39}$$

$$\epsilon_{21} = (-1)^{\wp_{21}} \Delta\left(\alpha_1, \alpha_2, \alpha_{12}\right) \left[\alpha_{45}, \alpha_{56}, \alpha_{123}, \alpha_{3456}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix}, \tag{A.40}$$

$$\wp_{21} = -\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 - \alpha_{12} - \alpha. \tag{A.41}$$

$$\epsilon_{22} = (-1)^{\wp_{22}} \Delta\left(\alpha_{123}, \alpha_{456}, \alpha\right) \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{56}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix}, \tag{A.42}$$

$$\wp_{22} = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_{123} + \alpha_{456}. \tag{A.43}$$

$$\begin{aligned} \epsilon_{23} &= (-1)^{\wp_{23}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{56}, \alpha_{456}, \alpha_{1234}\right]^{\frac{1}{2}} \\ &\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_{56} & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{1234} \end{Bmatrix}, \end{aligned} \tag{A.44}$$

$$\wp_{23} = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_5 + \alpha_6 - \alpha_{56} - \alpha + \alpha_{456}. \tag{A.45}$$

$$\begin{aligned} \epsilon_{24} &= (-1)^{\wp_{24}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{56}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{1234}\right]^{\frac{1}{2}} \\ &\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{Bmatrix} \begin{Bmatrix} \alpha_1 & \alpha_{23} & \alpha_{123} \\ \alpha_4 & \alpha_{1234} & \alpha_{234} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_{56} & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{1234} \end{Bmatrix}, \end{aligned} \tag{A.46}$$

$$\wp_{24} = \alpha_2 + \alpha_3 - \alpha_4 + \alpha_5 + \alpha_6 - \alpha_{23} - \alpha_{56} + \alpha_{456} - \alpha - \alpha_{1234}. \tag{A.47}$$

$$\epsilon_{25} = (-1)^{\wp_{25}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{56}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_1 & \alpha_{23} & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{23456} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_{56} & \alpha_{456} \\ \alpha_{23456} & \alpha_{23} & \alpha_{234} \end{array} \right\}, \tag{A.48}$$

$$\wp_{25} = \alpha_2 + \alpha_3 - \alpha_{123} + \alpha_5 + \alpha_6 - \alpha_{56} + \alpha + \alpha_{23456}. \tag{A.49}$$

$$\epsilon_{26} = (-1)^{\wp_{26}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{56}, \alpha_{123}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_1 & \alpha_{23} & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{23456} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{array} \right\}, \tag{A.50}$$

$$\wp_{26} = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 - \alpha_{23} + \alpha_{123} - \alpha. \tag{A.51}$$

$$\epsilon_{27} = (-1)^{\wp_{27}} \Delta\left(\alpha_1, \alpha_2, \alpha_{12}\right) \Delta\left(\alpha_{12}, \alpha_3, \alpha_{123}\right)$$

$$\times \left[\alpha_{45}, \alpha_{456}, \alpha_{1234}, \alpha_{12345}\right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{12345} & \alpha_{123} & \alpha_{1234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{array} \right\}, \tag{A.52}$$

$$\wp_{27} = -\alpha + \alpha_{12345} + \alpha_4 + \alpha_5 - \alpha_6 - \alpha_{45}. \tag{A.53}$$

$$\epsilon_{28} = (-1)^{\wp_{28}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{456}, \alpha_{1234}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{12345} & \alpha_{123} & \alpha_{1234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{array} \right\}, \tag{A.54}$$

$$\wp_{28} = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_{45} - \alpha_{123} + \alpha + \alpha_{12345}. \tag{A.55}$$

$$\epsilon_{29} = (-1)^{\wp_{29}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{1234}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_1 & \alpha_{23} & \alpha_{123} \\ \alpha_4 & \alpha_{1234} & \alpha_{234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{12345} & \alpha_{123} & \alpha_{1234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{array} \right\}, \tag{A.56}$$

$$\wp_{29} = \alpha_2 + \alpha_3 - \alpha_{23} + \alpha_{123} + \alpha_{1234} + \alpha_{12345} + \alpha - \alpha_5 + \alpha_6 - \alpha_{45}. \tag{A.57}$$

$$\epsilon_{30} = (-1)^{\wp_{30}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{2345}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_1 & \alpha_{23} & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{23456} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{2345} & \alpha_{23} & \alpha_{234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha_{23456} & \alpha_{23} & \alpha_{2345} \end{array} \right\}, \tag{A.58}$$

$$\wp_{30} = \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 - \alpha_{23} - \alpha_{45} - \alpha_{123} - \alpha_{2345} + \alpha + \alpha_{23456} + \alpha_{456}. \tag{A.59}$$

$$\epsilon_{31} = (-1)^{\wp_{31}} \left[\alpha_{12}, \alpha_{23}, \alpha_{45}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{2345}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_3 & \alpha_{123} & \alpha_{23} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_1 & \alpha_{23} & \alpha_{123} \\ \alpha_{45} & \alpha_{12345} & \alpha_{2345} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{2345} & \alpha_{23} & \alpha_{234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{array} \right\}, \tag{A.60}$$

$$\wp_{31} = \alpha_2 + \alpha_3 - \alpha_4 - \alpha_5 + \alpha_6 + \alpha_{2345} + \alpha + \alpha_{12345}. \tag{A.61}$$

$$\epsilon_{32} = (-1)^{\wp_{32}} \left[\alpha_{12}, \alpha_{34}, \alpha_{45}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{1234}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \left\{ \begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{34} & \alpha_{1234} & \alpha_{234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_4 & \alpha_{1234} & \alpha_{34} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{12345} & \alpha_{123} & \alpha_{1234} \end{array} \right\} \left\{ \begin{array}{ccc} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{array} \right\}, \tag{A.62}$$

$$\wp_{32} = \alpha_1 + \alpha_2 + \alpha_{34} + \alpha_6 + \alpha_{45} + \alpha + \alpha_5 + \alpha_{12345} + \alpha_3 + \alpha_{12}. \tag{A.63}$$

$$\epsilon_{33} = (-1)^{\wp_{33}} \left[\alpha_{12}, \alpha_{34}, \alpha_{45}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{2345}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \begin{bmatrix} \alpha_1 & & \alpha_2 & & \alpha_{234} & & \alpha_{2345} & & \alpha_{23456} & \\ & \alpha_{12} & & \alpha_{34} & & \alpha_5 & & \alpha_6 & & \alpha \\ \alpha_{123} & & \alpha_3 & & \alpha_4 & & \alpha_{45} & & \alpha_{456} & \end{bmatrix}, \tag{A.64}$$

$$\wp_{33} = \alpha_1 + \alpha_2 - \alpha_{12} + \alpha_{34} - \alpha_3 - \alpha_4. \tag{A.65}$$

$$\epsilon_{34} = (-1)^{\wp_{34}} \left[\alpha_{12}, \alpha_{34}, \alpha_{45}, \alpha_{123}, \alpha_{234}, \alpha_{456}, \alpha_{2345}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{Bmatrix} \begin{bmatrix} \alpha_2 & \alpha_{2345} & \alpha_3 & \alpha_{45} \\ \alpha_{12} & \alpha_{34} & \alpha_{12345} & \alpha_5 \\ \alpha_1 & \alpha_{123} & \alpha_{234} & \alpha_4 \end{bmatrix}, \tag{A.66}$$

$$\wp_{34} = -\alpha_{234} + \alpha_{2345} + 2\alpha_4 + \alpha + \alpha_6 + \alpha_{12} - \alpha_{34}. \tag{A.67}$$

$$\epsilon_{35} = (-1)^{\wp_{35}} \left[\alpha_{12}, \alpha_{34}, \alpha_{45}, \alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{3456}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{3456} & \alpha & \alpha_{23456} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{345} & \alpha_3 & \alpha_{34} \end{Bmatrix} \begin{Bmatrix} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha_{3456} & \alpha_3 & \alpha_{345} \end{Bmatrix}, \tag{A.68}$$

$$\wp_{35} = \alpha_{345} - \alpha_3 + \alpha_4 + \alpha_5 + \alpha_{45} + \alpha_6 + \alpha_{456} - \alpha_1 - \alpha_2 + \alpha_{12}. \tag{A.69}$$

$$\epsilon_{36} = (-1)^{\wp_{36}} \left[\alpha_{12}, \alpha_{34}, \alpha_{45}, \alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{2345}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{345} & \alpha_{12345} & \alpha_{2345} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{45} & \alpha_{12345} & \alpha_{345} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{345} & \alpha_3 & \alpha_{34} \end{Bmatrix} \begin{Bmatrix} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{Bmatrix}, \tag{A.70}$$

$$\wp_{36} = \alpha_1 + \alpha_2 - \alpha_4 - \alpha_5 + \alpha + \alpha_{12} - \alpha_{123} + \alpha_6. \tag{A.71}$$

$$\epsilon_{37} = (-1)^{\wp_{37}} \left[\alpha_{12}, \alpha_{34}, \alpha_{45}, \alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{2345}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_{345} & \alpha_3 & \alpha_{34} \end{Bmatrix} \begin{bmatrix} \alpha_2 & \alpha_{23456} & \alpha_3 & \alpha_{456} \\ \alpha_{12} & \alpha_{345} & \alpha & \alpha_6 \\ \alpha_1 & \alpha_{123} & \alpha_{2345} & \alpha_{45} \end{bmatrix}, \tag{A.72}$$

$$\wp_{37} = \alpha_4 + \alpha_5 + \alpha_{2345} - \alpha_{123} + \alpha_3 - \alpha_{23456} + \alpha_{456} + \alpha_{12}. \tag{A.73}$$

$$\epsilon_{38} = (-1)^{\wp_{38}} \Delta\left(\alpha_4, \alpha_5, \alpha_{45}\right) \left[\alpha_{12}, \alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{3456}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{3456} & \alpha & \alpha_{23456} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix} \begin{Bmatrix} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha_{3456} & \alpha_3 & \alpha_{345} \end{Bmatrix}, \tag{A.74}$$

$$\wp_{38} = \alpha_6 + \alpha_{45} - \alpha_{456} + 2\alpha - \alpha_1 - \alpha_2 - \alpha_{12}. \tag{A.75}$$

$$\epsilon_{39} = (-1)^{\wp_{39}} \Delta\left(\alpha_4, \alpha_5, \alpha_{45}\right) \left[\alpha_{12}, \alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{2345}, \alpha_{12345}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{345} & \alpha_{12345} & \alpha_{2345} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{45} & \alpha_{12345} & \alpha_{345} \end{Bmatrix} \begin{Bmatrix} \alpha_{45} & \alpha_6 & \alpha_{456} \\ \alpha & \alpha_{123} & \alpha_{12345} \end{Bmatrix}, \tag{A.76}$$

$$\wp_{39} = \alpha_1 + \alpha_2 + \alpha_{345} + \alpha - \alpha_{12} + \alpha_{123} - \alpha_3 + \alpha_6. \tag{A.77}$$

$$\epsilon_{40} = (-1)^{\wp_{40}} \Delta\left(\alpha_4, \alpha_5, \alpha_{45}\right) \left[\alpha_{12}, \alpha_{123}, \alpha_{345}, \alpha_{456}, \alpha_{2345}, \alpha_{23456}\right]^{\frac{1}{2}} \begin{bmatrix} \alpha_1 & \alpha_{2345} & \alpha_{123} & \alpha_{45} \\ \alpha_{12} & \alpha & \alpha_{345} & \alpha_6 \\ \alpha_2 & \alpha_3 & \alpha_{23456} & \alpha_{456} \end{bmatrix}, \tag{A.78}$$

$$\wp_{40} = \alpha_{123} + \alpha_{2345} + 2\alpha_2 + \alpha_{23456} - \alpha_{456} - \alpha_{12} - \alpha_{345}. \tag{A.79}$$

$$\epsilon_{41} = (-1)^{\wp_{41}} \Delta\left(\alpha_4, \alpha_5, \alpha_{45}\right) \Delta\left(\alpha_{45}, \alpha_6, \alpha_{456}\right)$$

$$\times \left[\alpha_{12}, \alpha_{123}, \alpha_{3456}, \alpha_{23456}\right]^{\frac{1}{2}} \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{3456} & \alpha & \alpha_{23456} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix}, \tag{A.80}$$

$$\wp_{41} = \alpha_{12} - \alpha_1 - \alpha_2 + \alpha_3 + \alpha_{456} - \alpha_{3456}. \tag{A.81}$$

$$\epsilon_{42} = (-1)^{\wp_{42}} \left[\alpha_{12}, \alpha_{45}, \alpha_{56}, \alpha_{123}, \alpha_{3456}, \alpha_{23456}\right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} \alpha_1 & \alpha_2 & \alpha_{12} \\ \alpha_{3456} & \alpha & \alpha_{23456} \end{Bmatrix} \begin{Bmatrix} \alpha_{12} & \alpha_3 & \alpha_{123} \\ \alpha_{456} & \alpha & \alpha_{3456} \end{Bmatrix} \begin{Bmatrix} \alpha_4 & \alpha_5 & \alpha_{45} \\ \alpha_6 & \alpha_{456} & \alpha_{56} \end{Bmatrix}, \tag{A.82}$$

$$\wp_{42} = -\alpha_4 - \alpha_5 - \alpha_6 + \alpha_{12} - \alpha_1 - \alpha_2 + \alpha_3 - \alpha_{3456}. \tag{A.83}$$

The quantities $\{\ldots\}$ denote $6j$-coefficients of SU (2). The definition of $12j$-coefficient of the second kind (see Eqs. (A.23), (A.66), (A.72), (A.78)) one can find in [16, 17]. The definition of $15j$-coefficient of the second kind (see Eq. (A.64)) is presented in [16, 18, 19].

B Classes

In this Appendix the data, required for the study of quantities in Sec. 5, are presented. In tables operators $\langle x_\pi \rangle$, pertaining to the specific class $X_n(\Delta_1, \Delta_2, \ldots, \Delta_n)$ with $n \in \{2, 3, \ldots, 6\}$, are listed. The explicit expressions for $\widehat{\pi}$ (see Eq. (5.4)) and $\widehat{\widehat{\pi}}$ (see Eqs. (5.12)-(5.13)) are given.

$X_2\,(0,0)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle 112112 \rangle$	(35)	$\langle 111122 \rangle$
$\langle 121112 \rangle$	(25)	
$\langle 211112 \rangle$	(15)	
$\langle 121121 \rangle$	$-\,(26)$	
$\langle 112121 \rangle$	(36)	
$\langle 112211 \rangle$	$(35)\,(46)$	
$\langle 122122 \rangle$	$-\,(24)$	$\langle 112222 \rangle$
$\langle 122212 \rangle$	(25)	
$\langle 122221 \rangle$	$-\,(26)$	
$\langle 212122 \rangle$	(14)	
$\langle 212212 \rangle$	$-\,(15)$	
$\langle 221122 \rangle$	$(13)\,(24)$	

Tab. 8: Operators of class $X_2\,(0,0)$.

$X_2\,(+1,-1)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle 111112 \rangle$	1_6	$\langle 111112 \rangle$
$\langle 111121 \rangle$	(56)	
$\langle 111211 \rangle$	$-\,(46)$	
$\langle 211122 \rangle$	(14)	$\langle 111222 \rangle$
$\langle 121122 \rangle$	$-\,(24)$	
$\langle 112122 \rangle$	(34)	
$\langle 211212 \rangle$	$-\,(15)$	
$\langle 121212 \rangle$	(25)	
$\langle 112212 \rangle$	$-\,(35)$	
$\langle 211221 \rangle$	(16)	
$\langle 121221 \rangle$	$-\,(26)$	
$\langle 112221 \rangle$	(36)	
$\langle 122222 \rangle$	1_6	$\langle 122222 \rangle$
$\langle 212222 \rangle$	(12)	
$\langle 221222 \rangle$	$-\,(13)$	

Tab. 9: Operators of class $X_2\,(+1,-1)$.

$X_2\,(+2,-2)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle 111122 \rangle$	1_6	$\langle 111122 \rangle$
$\langle 111212 \rangle$	(45)	
$\langle 111221 \rangle$	$-\,(46)$	
$\langle 112222 \rangle$	1_6	$\langle 112222 \rangle$
$\langle 121222 \rangle$	(23)	
$\langle 211222 \rangle$	$-\,(13)$	

Tab. 10: Operators of class $X_2\,(+2,-2)$.

The class $X_2\,(+3,-3)$ is of order $d = 1$. The only operator which belongs to $X_2\,(+3,-3)$, is $\langle 111222 \rangle$.

$X_3\,(0,0,0)$			$X_3\,(0,0,0)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$	$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle 123\,\{123\} \rangle$	$-\,(24)\,(35)\,\widehat{\vartheta}$	$\langle 112233 \rangle$	$\langle 213213 \rangle$	$-\,(135)$	$\langle 112233 \rangle$
$\langle 132123 \rangle$	(254)		$\langle 213312 \rangle$	$-\,(135)\,(46)$	
$\langle 132132 \rangle$	$-\,(264)$		$\langle 231123 \rangle$	$(13)\,(254)$	
$\langle 132213 \rangle$	(25)		$\langle 231132 \rangle$	$-\,(13)\,(264)$	
$\langle 132231 \rangle$	$-\,(26)$		$\langle 231213 \rangle$	$(13)\,(25)$	
$\langle 132312 \rangle$	$(25)\,(46)$		$\langle 312123 \rangle$	$-\,(154)$	
$\langle 213123 \rangle$	$(14)\,(35)$		$\langle 312213 \rangle$	$-\,(15)$	
$\langle 213132 \rangle$	$-\,(14)\,(36)$		$\langle 321123 \rangle$	$-\,(154)\,(23)$	

Tab. 11: Operators of class $X_3\,(0,0,0)$.

$X_3\,(+2,-1,-1)$			$X_3\,(+2,-1,-1)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$	$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 111\,\{123\}\rangle$	$\widehat{\vartheta}$	$\langle 111123\rangle$	$\langle 113233\rangle$	(34)	$\langle 112333\rangle$
$\langle 112223\rangle$	1_6	$\langle 112223\rangle$	$\langle 113323\rangle$	$-\,(35)$	
$\langle 112232\rangle$	(56)		$\langle 113332\rangle$	(36)	
$\langle 112322\rangle$	$-\,(46)$		$\langle 131233\rangle$	(243)	
$\langle 121223\rangle$	(23)		$\langle 131323\rangle$	$-\,(253)$	
$\langle 121232\rangle$	$(23)\,(56)$		$\langle 131332\rangle$	(263)	
$\langle 121322\rangle$	$-\,(23)\,(46)$		$\langle 311233\rangle$	$-\,(143)$	
$\langle 211223\rangle$	$-\,(13)$		$\langle 311323\rangle$	(153)	
$\langle 211232\rangle$	$-\,(13)\,(56)$		$\langle 311332\rangle$	$-\,(163)$	
$\langle 211322\rangle$	$(13)\,(46)$				
$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$			$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$		
$X_3\,(-1,+2,-1)$			$X_3\,(-1,-1,+2)$		
$\langle x\rangle$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$	$\langle x\rangle$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$\langle 111123\rangle$	$-\,(15)$	$\langle 122223\rangle$	$\langle 111123\rangle$	$-\,(16)\,(25)$	$\langle 123333\rangle$
$\langle 112223\rangle$	$(14)\,(25)$	$\langle 111223\rangle$	$\langle 112223\rangle$	$-\,(16)\,(25)$	$\langle 122233\rangle$
$\langle 112333\rangle$	$-\,(13)$	$\langle 122333\rangle$	$\langle 112333\rangle$	$(16)\,(25)$	$\langle 111233\rangle$

Tab. 12: Operators of classes $X_3\,(\Delta_1,\Delta_2,\Delta_3)$ with $\Delta_x = -1, +1, +2$.

$X_3\,(+3,-2,-1)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 111223\rangle$	1_6	$\langle 111223\rangle$
$\langle 111232\rangle$	(56)	
$\langle 111322\rangle$	$-\,(46)$	
$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$X_3\,(-2,+3,-1)$	$(15)\,(24)$	$\langle 112223\rangle$
$X_3\,(-1,-2,+3)$	$(16)\,(24)\,(35)$	$\langle 122333\rangle$
$X_3\,(+3,-1,-2)$	$-\,(46)$	$\langle 111233\rangle$
$X_3\,(-1,+3,-2)$	(146)	$\langle 122233\rangle$
$X_3\,(-2,-1,+3)$	$(14)\,(25)\,(36)$	$\langle 112333\rangle$

Tab. 13: Operators of classes $X_3\,(\Delta_1,\Delta_2,\Delta_3)$ with $\Delta_x = -2, -1, +3$.

$X_3\,(+1,-1,0)$			$X_3\,(+1,-1,0)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$	$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 133233\rangle$	$-\,(24)$	$\langle 123333\rangle$	$\langle 313332\rangle$	$-\,(162)$	$\langle 123333\rangle$
$\langle 133323\rangle$	(25)		$\langle 331233\rangle$	$(13)\,(24)$	
$\langle 133332\rangle$	$-\,(26)$		$\langle 331323\rangle$	$-\,(13)\,(25)$	
$\langle 313233\rangle$	$-\,(142)$		$\langle 331332\rangle$	$(13)\,(26)$	
$\langle 313323\rangle$	(152)				
$\langle 113\,\{123\}\rangle$	$(354)\,\widehat{\vartheta}$	$\langle 111233\rangle$	$\langle \{123\}\,223\rangle$	$-\,(35)\,\widehat{\eta}$	$\langle 122233\rangle$
$\langle 131\,\{123\}\rangle$	$-\,(254)\,\widehat{\vartheta}$		$\langle \{123\}\,232\rangle$	$(36)\,\widehat{\eta}$	
$\langle 311\,\{123\}\rangle$	$(154)\,\widehat{\vartheta}$		$\langle \{123\}\,322\rangle$	$(35)\,(46)\,\widehat{\eta}$	
$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$			$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$		
$X_3\,(+1,0,-1)$			$X_3\,(0,+1,-1)$		
$\langle x\rangle$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$	$\langle x\rangle$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$\langle 123333\rangle$	$-\,(162)$	$\langle 122223\rangle$	$\langle 123333\rangle$	$(15)\,(26)$	$\langle 111123\rangle$
$\langle 111233\rangle$	$-\,(46)$	$\langle 111223\rangle$	$\langle 111233\rangle$	$(15)\,(246)$	$\langle 112223\rangle$
$\langle 122233\rangle$	$(25)\,(36)$	$\langle 122333\rangle$	$\langle 122233\rangle$	$-\,(135)\,(26)$	$\langle 112333\rangle$

Tab. 14: Operators of classes $X_3\,(\Delta_1,\Delta_2,\Delta_3)$ with $\Delta_x = -1, 0, +1$.

$X_3\,(+2,-2,0)$			$X_3\,(+2,-2,0)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$	$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 113223\rangle$	$-\,(35)$	$\langle 112233\rangle$	$\langle 131322\rangle$	$(253)\,(46)$	$\langle 112233\rangle$
$\langle 113232\rangle$	(36)		$\langle 311223\rangle$	(153)	
$\langle 113322\rangle$	$(35)\,(46)$		$\langle 311232\rangle$	$-\,(163)$	
$\langle 131223\rangle$	$-\,(253)$		$\langle 311322\rangle$	$-\,(153)\,(46)$	
$\langle 131232\rangle$	(263)				
$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$	$X_3\,(\Delta'_1,\Delta'_2,\Delta'_3)$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$X_3\,(+2,0,-2)$	$(35)\,(46)$	$\langle 112233\rangle$	$X_3\,(0,+2,-2)$	$(135)\,(246)$	$\langle 112233\rangle$

Tab. 15: Operators of classes $X_3\,(\Delta_1,\Delta_2,\Delta_3)$ with $\Delta_x=-2,0,+2$.

$X_4\,(+1,+1,-1,-1)$			$X_4\,(+1,+1,-1,-1)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$	$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 112\,\{134\}\rangle$	$(34)\,\widehat{\vartheta}$	$\langle 111234\rangle$	$\langle 122\,\{234\}\rangle$	$\widehat{\vartheta}$	$\langle 122234\rangle$
$\langle 121\,\{134\}\rangle$	$-\,(24)\,\widehat{\vartheta}$		$\langle 212\,\{234\}\rangle$	$(12)\,\widehat{\vartheta}$	
$\langle 211\,\{134\}\rangle$	$(14)\,\widehat{\vartheta}$		$\langle 221\,\{234\}\rangle$	$-\,(13)\,\widehat{\vartheta}$	
$\langle \{123\}\,334\rangle$	$\widehat{\eta}$	$\langle 123334\rangle$	$\langle \{124\}\,344\rangle$	$(34)\,\widehat{\eta}$	$\langle 123444\rangle$
$\langle \{123\}\,343\rangle$	$(56)\,\widehat{\eta}$		$\langle \{124\}\,434\rangle$	$-\,(35)\,\widehat{\eta}$	
$\langle \{123\}\,433\rangle$	$-\,(46)\,\widehat{\eta}$		$\langle \{124\}\,443\rangle$	$(36)\,\widehat{\eta}$	
$X_4\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4)$			$X_4\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4)$		
$X_4\,(+1,-1,+1,-1)$			$X_4\,(+1,-1,-1,+1)$		
$\langle x\rangle$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$	$\langle x\rangle$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$\langle 111234\rangle$	(45)	$\langle 111234\rangle$	$\langle 111234\rangle$	(46)	$\langle 111234\rangle$
$\langle 122234\rangle$	(25)	$\langle 123334\rangle$	$\langle 122234\rangle$	$-\,(26)\,(35)$	$\langle 123444\rangle$
$\langle 123334\rangle$	(25)	$\langle 122234\rangle$	$\langle 123334\rangle$	(26)	$\langle 123334\rangle$
$\langle 123444\rangle$	(23)	$\langle 123444\rangle$	$\langle 123444\rangle$	$-\,(26)\,(35)$	$\langle 122234\rangle$

Tab. 16: Operators of classes $X_4\,(\Delta_1,\Delta_2,\Delta_3,\Delta_4)$ with $\Delta_x=-1,+1$.

$X_4\,(+2,-2,+1,-1)$			$X_4\,(+2,-2,+1,-1)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$	$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 113224\rangle$	$-\,(35)$	$\langle 112234\rangle$	$\langle 131422\rangle$	$(253)\,(46)$	$\langle 112234\rangle$
$\langle 113242\rangle$	$-\,(356)$		$\langle 311224\rangle$	$-\,(1532)$	
$\langle 113422\rangle$	$(35)\,(46)$		$\langle 311242\rangle$	(1563)	
$\langle 131224\rangle$	$-\,(253)$		$\langle 311422\rangle$	$-\,(153)\,(46)$	
$\langle 131242\rangle$	$-\,(2563)$				
$X_4\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4)$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$	$X_4\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4)$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$X_4\,(+2,-2,-1,+1)$	(56)	$\langle 112234\rangle$	$X_4\,(+1,+2,-2,-1)$	(135)	$\langle 122334\rangle$
$X_4\,(+2,+1,-2,-1)$	$-\,(35)$	$\langle 112334\rangle$	$X_4\,(+1,+2,-1,-2)$	$-\,(135)\,(46)$	$\langle 122344\rangle$
$X_4\,(+2,+1,-1,-2)$	$(35)\,(46)$	$\langle 112344\rangle$	$X_4\,(+1,-2,+2,-1)$	$(15)\,(24)$	$\langle 122334\rangle$
$X_4\,(+2,-1,-2,+1)$	$-\,(356)$	$\langle 112334\rangle$	$X_4\,(+1,-2,-1,+2)$	$-\,(15)\,(264)$	$\langle 122344\rangle$
$X_4\,(+2,-1,+1,-2)$	$-\,(36)\,(45)$	$\langle 112344\rangle$	$X_4\,(+1,-1,+2,-2)$	$(135)\,(246)$	$\langle 123344\rangle$
			$X_4\,(+1,-1,-2,+2)$	$(15)\,(26)$	$\langle 123344\rangle$

Tab. 17: Operators of classes $X_4\,(\Delta_1,\Delta_2,\Delta_3,\Delta_4)$ with $\Delta_x=-2,-1,+1,+2$.

$X_4\,(+3,-1,-1,-1)$		
$\langle x_\pi\rangle$	$\widehat{\pi}$	$\langle x\rangle$
$\langle 111\,\{234\}\rangle$	$\widehat{\vartheta}$	$\langle 111234\rangle$
$X_4\,(\Delta'_1,\Delta'_2,\Delta'_3,\Delta'_4)$	$\widehat{\widetilde{\pi}}$	$\langle y\rangle$
$X_4\,(-1,+3,-1,-1)$	(14)	$\langle 122234\rangle$
$X_4\,(-1,-1,+3,-1)$	$-\,(15)\,(24)$	$\langle 123334\rangle$
$X_4\,(-1,-1,-1,+3)$	$(16)\,(24)\,(35)$	$\langle 123444\rangle$

Tab. 18: Operators of classes $X_4\,(\Delta_1,\Delta_2,\Delta_3,\Delta_4)$ with $\Delta_x=-1,+3$.

$X_4\,(+1,-1,0,0)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle \{134\}\,\{234\} \rangle$	$-\,(24)\,(35)\,\widehat{\eta}\widehat{\vartheta}$	$\langle 123344 \rangle$
$X_4\,(\Delta_1', \Delta_2', \Delta_3', \Delta_4')$	$\widehat{\tilde{\pi}}$	$\langle y \rangle$
$X_4\,(+1,0,-1,0)$	$-\,(24)$	$\langle 122344 \rangle$
$X_4\,(+1,0,0,-1)$	$(26)\,(35)$	$\langle 122334 \rangle$
$X_4\,(0,+1,-1,0)$	$(13)\,(24)$	$\langle 112344 \rangle$
$X_4\,(0,+1,0,-1)$	$-\,(135)\,(26)$	$\langle 112334 \rangle$
$X_4\,(0,0,+1,-1)$	$(15)\,(26)$	$\langle 112234 \rangle$

Tab. 19: Operators of classes $X_4\,(\Delta_1, \Delta_2, \Delta_3, \Delta_4)$ with $\Delta_x = -1, 0, +1$.

$X_4\,(+2,-1,-1,0)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle 114\,\{234\} \rangle$	$(354)\,\widehat{\vartheta}$	$\langle 112344 \rangle$
$\langle 141\,\{234\} \rangle$	$(2543)\,\widehat{\vartheta}$	
$\langle 411\,\{234\} \rangle$	$-\,(1543)\,\widehat{\vartheta}$	
$X_4\,(\Delta_1', \Delta_2', \Delta_3', \Delta_4')$	$\widehat{\tilde{\pi}}$	$\langle y \rangle$
$X_4\,(-1,+2,-1,0)$	$-\,(13)$	$\langle 122344 \rangle$
$X_4\,(+2,-1,0,-1)$	$-\,(46)$	$\langle 112334 \rangle$
$X_4\,(+2,0,-1,-1)$	$-\,(36)\,(45)$	$\langle 112234 \rangle$
$X_4\,(0,+2,-1,-1)$	$-\,(145)\,(236)$	$\langle 112234 \rangle$
$X_4\,(0,-1,+2,-1)$	$-\,(15)\,(2436)$	$\langle 112334 \rangle$
$X_4\,(-1,+2,0,-1)$	(1364)	$\langle 122334 \rangle$
$X_4\,(-1,-1,+2,0)$	$(13)\,(24)$	$\langle 123344 \rangle$
$X_4\,(-1,-1,0,+2)$	$-\,(164)\,(253)$	$\langle 123344 \rangle$
$X_4\,(-1,0,+2,-1)$	$(14)\,(25)\,(36)$	$\langle 122334 \rangle$
$X_4\,(-1,0,-1,+2)$	$-\,(1634)\,(25)$	$\langle 122344 \rangle$
$X_4\,(0,-1,-1,+2)$	$(15)\,(26)$	$\langle 112344 \rangle$

Tab. 20: Operators of classes $X_4\,(\Delta_1, \Delta_2, \Delta_3, \Delta_4)$ with $\Delta_x = -1, 0, +2$.

$X_5\,(+2,+1,-1,-1,-1)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle 112\,\{345\} \rangle$	$\widehat{\vartheta}$	$\langle 112345 \rangle$
$\langle 121\,\{345\} \rangle$	$(23)\,\widehat{\vartheta}$	
$\langle 211\,\{345\} \rangle$	$-\,(13)\,\widehat{\vartheta}$	
$X_5\,(\Delta_1', \Delta_2', \Delta_3', \Delta_4', \Delta_5')$	$\widehat{\tilde{\pi}}$	$\langle y \rangle$
$X_5\,(+2,-1,+1,-1,-1)$	(34)	$\langle 112345 \rangle$
$X_5\,(+2,-1,-1,+1,-1)$	(35)	
$X_5\,(+2,-1,-1,-1,+1)$	(36)	
$X_5\,(+1,+2,-1,-1,-1)$	$-\,(13)$	$\langle 122345 \rangle$
$X_5\,(-1,+2,+1,-1,-1)$	$-\,(134)$	
$X_5\,(-1,+2,-1,+1,-1)$	$-\,(135)$	
$X_5\,(-1,+2,-1,-1,+1)$	$-\,(136)$	
$X_5\,(-1,-1,+2,+1,-1)$	$(135)\,(24)$	$\langle 123345 \rangle$
$X_5\,(-1,+1,+2,-1,-1)$	$-\,(14)\,(23)$	
$X_5\,(+1,-1,+2,-1,-1)$	$(13)\,(24)$	
$X_5\,(-1,-1,+2,-1,+1)$	$(136)\,(24)$	
$X_5\,(-1,+1,-1,+2,-1)$	$(14)\,(253)$	$\langle 123445 \rangle$
$X_5\,(+1,-1,-1,+2,-1)$	$-\,(153)\,(24)$	
$X_5\,(-1,-1,+1,+2,-1)$	$-\,(15)\,(24)$	
$X_5\,(-1,-1,-1,+2,+1)$	$(1436)\,(25)$	
$X_5\,(+1,-1,-1,-1,+2)$	$(1543)\,(26)$	$\langle 123455 \rangle$
$X_5\,(-1,-1,+1,-1,+2)$	$-\,(16)\,(254)$	
$X_5\,(-1,+1,-1,-1,+2)$	$(154)\,(263)$	
$X_5\,(-1,-1,-1,+1,+2)$	$(15)\,(2634)$	

Tab. 21: Operators of classes $X_5\,(\Delta_1, \Delta_2, \Delta_3, \Delta_4, \Delta_5)$ with $\Delta_x = -1, +1, +2$.

$X_5\,(+1, +1, -1, -1, 0)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle \{125\}\,\{345\} \rangle$	$(354)\,\widehat{\eta}\widehat{\vartheta}$	$\langle 123455 \rangle$
$X_5\,(\Delta'_1, \Delta'_2, \Delta'_3, \Delta'_4, \Delta'_5)$	$\widehat{\widehat{\pi}}$	$\langle y \rangle$
$X_5\,(+1, -1, +1, -1, 0)$	(23)	$\langle 123455 \rangle$
$X_5\,(+1, -1, -1, +1, 0)$	(24)	
$X_5\,(+1, +1, -1, 0, -1)$	$-\,(46)$	$\langle 123445 \rangle$
$X_5\,(+1, -1, +1, 0, -1)$	$-\,(23)\,(46)$	
$X_5\,(+1, -1, -1, 0, +1)$	$-\,(264)$	
$X_5\,(+1, +1, 0, -1, -1)$	$(35)\,(46)$	$\langle 123345 \rangle$
$X_5\,(+1, -1, 0, +1, -1)$	$-\,(254)\,(36)$	
$X_5\,(+1, -1, 0, -1, +1)$	$-\,(263)\,(45)$	
$X_5\,(+1, 0, +1, -1, -1)$	$(245)\,(36)$	$\langle 122345 \rangle$
$X_5\,(+1, 0, -1, +1, -1)$	$(25)\,(346)$	
$X_5\,(+1, 0, -1, -1, +1)$	$-\,(26)\,(345)$	
$X_5\,(0, +1, +1, -1, -1)$	$-\,(146)\,(235)$	$\langle 112345 \rangle$
$X_5\,(0, +1, -1, +1, -1)$	$(15)\,(2346)$	
$X_5\,(0, +1, -1, -1, +1)$	$-\,(16)\,(2345)$	

Tab. 22: Operators of classes $X_5\,(\Delta_1, \Delta_2, \Delta_3, \Delta_4, \Delta_5)$ with $\Delta_x = -1, 0, +1$.

$X_6\,(+1, +1, +1, -1, -1, -1)$		
$\langle x_\pi \rangle$	$\widehat{\pi}$	$\langle x \rangle$
$\langle \{123\}\,\{456\} \rangle$	$\widehat{\eta}\widehat{\vartheta}$	$\langle 123456 \rangle$
$X_6\,(\Delta'_1, \Delta'_2, \Delta'_3, \Delta'_4, \Delta'_5, \Delta'_6)$	$\widehat{\widehat{\pi}}$	$\langle y \rangle$
$X_6\,(+1, +1, -1, +1, -1, -1)$	(34)	$\langle 123456 \rangle$
$X_6\,(+1, +1, -1, -1, +1, -1)$	(35)	
$X_6\,(+1, +1, -1, -1, -1, +1)$	(36)	
$X_6\,(+1, -1, +1, +1, -1, -1)$	(24)	
$X_6\,(+1, -1, +1, -1, +1, -1)$	(254)	
$X_6\,(+1, -1, +1, -1, -1, +1)$	(264)	
$X_6\,(+1, -1, -1, +1, +1, -1)$	$(24)\,(35)$	
$X_6\,(+1, -1, -1, +1, -1, +1)$	$(24)\,(36)$	
$X_6\,(+1, -1, -1, -1, +1, +1)$	$(25)\,(36)$	

Tab. 23: Operators of classes $X_6\,(\Delta_1, \Delta_2, \Delta_3, \Delta_4, \Delta_5, \Delta_6)$ with $\Delta_x = -1, +1$.

References

[1] B. Brandow, Rev. Mod. Phys. 39 (1967) 771

[2] V. Kvasnička, Chem. J. Phys. B 24 (1974) 605

[3] I. Lindgren, J. Phys. B 7 (1974) 2441

[4] I. Lindgren and J. Morrison, Atomic Many-Body Theory, Springer Series in Chemical Physics, Vol. 13 (1982)

[5] R. Karazija, Introduction to the Theory of X-ray and Electronic Spectra of Free Atoms, Plenum Press, New York, (1996)

[6] G. Merkelis, R. Karazija, J. Electron. Spectr. Relat. Phenom. 133 (2003) 123

[7] Brian R. Judd, Second Quantization and Atomic Spectroscopy, The Johns Hopkins Press, Baltimore (1967)

[8] B. R. Judd, Operator Techniques in Atomic Spectroscopy, McGraw-Hill, New York, (1963)

[9] Z. Rudzikas and J. Kaniauskas, Quasispin and Isospin in the Theory of Atom, Mokslas, Vilnius (1984)

[10] Zenonas Rudzikas, Theoretical Atomic Spectroscopy, Cambridge University Press, Cambridge (1997)

[11] G. Merkelis, Physica Scripta 61 (2000) 662

[12] G. Gaigalas, Z. Rudzikas, Ch. F. Fischer, J. Phys. B: At. Mol. Opt. Phys. 30 (1997) 3747-3771

[13] G. Gaigalas, S. Fritzsche, I. P. Grant, Comp. Phys. Commun. 139 (2001) 263-278

[14] R. Juršėnas and G. Merkelis, Cent. Eur. J. Phys., DOI:10.2478/s11534-009-0126-5 (2009)

[15] R. Juršėnas, G. Merkelis, At. Data Nucl. Data Tables (accepted, 2009)

[16] A. Jucys, Y. Levinson, V. Vanagas, Mahematical Apparatus of the Theory of Angular Momentum, New York (1964)

[17] V. V. Vanagas and J. V. Čiplys, Trudy AN Lit. SSR, Ser. B 3 (15) (1958) 17

[18] S. Ališauskas, arXiv:math/9912142 (2002)

[19] S. Ališauskas, arXiv:math-ph/0509035 (2004)